Homework #2

Due: April 04, 2018 (Wed), in class

Instructor: Yongju Choi

1. You drank three bottles of soju last night. You want to know how much ethanol you drank and how it can be translated into calories. Assume that soju is a pure 17.8%¹ ethanol solution in water (volume ethanol/volume solution) and also that when ethanol is mixed with water, the total volume remains unchanged (for example, 1 L soju contains 178 mL ethanol and 822 mL water). Using the following values, answer the questions.

```
density of ethanol = 0.78 kg/L
density of water = 1.00 kg/L
```

¹참이슬 후레쉬 도수

- 1) Calculate the mass concentration (in g/L), molarity (in M) and mole fraction of ethanol in soju. (10 points)
- 2) Estimate the mass of ethanol you drank (in grams) and the energy obtained by the intake of ethanol (in kcal). The enthalpy of reaction for combustion of ethanol is -295 kcal/mole. This means that by 1 mole of ethanol you can get 295 kcal of energy (at maximum). A bottle of soju is 360 mL in volume.

(Note: The lethal dose of ethanol is approximately 6 g ethanol/kg body weight. This means for a 60 kg person, 360 g of ethanol can kill. You may get some guide to protect yourselves from drinking soju by the calculation.)

(10 points)

2. The BOD first-order reaction rate constant for a water sample is determined to be 0.25/d at 20° C. Using the modified van't Hoff-Arrhenius relationship, i) determine the rate constants at 4° C, 14° C and 24° C and ii) calculate the time required to degrade 50% of the UBOD in the sample at the three temperatures. The temperature coefficient θ is given as 1.135 for $4-20^{\circ}$ C and 1.056 for $20-30^{\circ}$ C.

(15 points)

3. Using the following data for water sample analysis, answer the followings.

Instructor: Yongju Choi

Ions	Concentration (mg/L)
Ca ²⁺	76.0
Mg ²⁺ Na ⁺	26.8
Na ⁺	23.0
K ⁺	19.6
Cl	37.2
SO ₄ ²⁻ CO ₃ ²⁻	192.0
CO_3^{2-}	2.7
HCO ₃ -	136.5

1) Determine the acceptance of the analysis. (10 points)

2) Determine the sodium adsorption ratio (SAR). In terms of the SAR, is the water acceptable for irrigation? (5 points)

3) Estimate the pH of the sample. Use for pK_{a2} value of 10.3 for the dissociation of HCO_3^- into CO_3^{2-} relationship. (6 points)

4) Determine the alkalinity, total hardness and carbonate hardness in mg/L as CaCO₃. (9 points)

4. 물 속에 존재하는 유기물질을 화학식 $C_{10}H_{19}O_3N$ 으로 표현할 수 있다고 가정하자. 환경 정책기본법 시행령 상 하천의 생활환경기준 중 TOC 기준으로 IV등급을 겨우 만족하는 물은 IV등급의 COD 기준을 만족하겠는가? 다음의 화학식을 이용하시오.

$$C_{10}H_{19}O_3N + 12.5O_2 \rightarrow 9CO_2 + 7H_2O + HCO_3^- + NH_4^+$$

(15 points)

(참조: 법령정보는 http://www.law.go.kr에서 찾아볼 수 있음)

5. 10⁻³ M acetic acid (CH₃COOH; HAc) is dissolved in pure water. For this solution, answer the followings.

Instructor: Yongju Choi

- 1) Draw the pH-pC diagram for 10⁻³ HAc and mark an intersect of the plot that satisfies the charge balance. Estimate the pH of this solution from the intersect. (5 points)
- 2) In the acetic acid solution, 10⁻⁵ M sodium hypochlorite (NaOCl) is added. On the 10⁻³ HAc pH-pC diagram, plot pC for HOCl and OCl at the C_T of 10⁻⁵ M. Mark the intersection of the pH-pC diagram that satisfies the mass balance and charge balance. Estimate the pH of this solution from the intersect. How does the solution pH change by the addition of NaOCl? (10 points)
- 3) From the pH-pC diagram, estimate the concentrations of HOCl and OCl in the solution. (5 points)
- 6. A pH buffer is made using a weak acid and a salt of a conjugate base of the weak acid. For a pH buffer made using a weak monoprotic acid (represented as HA) system, develop a program that can calculate the concentrations of the weak acid (HA; with any pK_a value) and the salt (MA; M represents a well-ionizing metal (i.e., perfectly dissolved in water to be M⁺) such as Na and K) that should be added to make a pH buffer solution of any pH and C_T value. You can use any program including Microsoft Excel and MATLAB.

(50 points)