





Writing half reactions – example

example 1) Glucose oxidation








in redox reactions, water is almost always a reactant or a product


because we write half reactions in the direction of reduction, e^- always appear on the left



example 2) Nitrate reduction



Textbook example: Alanine oxidation



Eq. [2.22] to [2.25] in the textbook: slightly different equations depending on different forms of carbonate species and ammonia, and different source of hydrogen (all are correct)

Ammonia $\text{pK}_a = 9.25 \rightarrow$ mostly in NH_4^+ in neutral pH

Carbonate $\text{pK}_{a1} = 6.3$, $\text{pK}_{a2} = 10.3 \rightarrow$ mostly in HCO_3^- with some in H_2CO_3 in neutral pH

Thus, some find it better to use NH_4^+ and HCO_3^- to write reactions, but some prefer others for different reasons




Overall reactions for bacterial growth

We can write up overall reactions for bacterial growth, which involves reactions for energy production and cell synthesis → full stoichiometry of what's happening when you grow bacteria in a growth medium!

- Combining electron donor and acceptor half reactions, we get the overall reaction for respiration (energy reaction)

$$R_e = R_a - R_d \quad (- \text{ sign on } R_d \text{ because } R_d \text{ is written in the direction of reduction})$$



ex) Benzoate respiration using nitrate as an electron acceptor (textbook)



- Combining cell synthesis and electron donor half reactions, we get the overall reaction for cell synthesis using a specific electron donor (synthesis reaction)

$$R_s = R_c - R_d$$

ex) Use of benzoate to synthesize cell with generic cell formula (textbook)



- Now, note that for the overall reactions for bacterial growth, R_e occurs with a fraction of f_e and R_s occurs with a fraction of f_s . Therefore,


$$R = f_e R_e + f_s R_s \quad [\text{Eq (1)}]$$

ex) Overall cell synthesis reaction for benzoate as an electron donor and nitrate as an electron acceptor (textbook)

when $f_s = 0.40$,

$$f_e = 1 - f_s = 0.60$$

$$R = 0.60 \times R_e + 0.40 \times R_s$$



→ Full stoichiometry obtained!

If you plug in the expressions for R_e and R_s in Eq (1),

$$R = f_e(R_a - R_d) + f_s(R_c - R_d) = f_e R_a + f_s R_c - R_d$$

So, instead of calculating the energy and cell synthesis reactions separately, you can use the equation above to make the overall reaction.