

Ch. 6. Quantum Theory of the H Atom



Erwin Schrödinger
(1887-1961)


... Schrödinger gave a talk on de Broglie's notion that a moving particle has a wave character. A colleague remarked to him afterward that to deal properly with a wave, one needs a wave equation. Schrödinger took this to heart, and a few weeks later he was "struggling with a new atomic theory. If only I knew more mathematics! I am very optimistic about this thing and expect that if I can only ... solve it, it will be *very* beautiful."

The struggle was successful, and in January 1926 the first of four papers on "Quantization as an Eigenvalue Problem" was completed. In this epochal paper Schrödinger introduced the equation that bears his name and solved it for the hydrogen atom, thereby opening wide the door to the modern view of the atom which others had only pushed ajar.
-- A. Beiser





Spherical Coordinates



© 2005 Brooks/Cole - Thomson





When Potential Depends on r Only

Time-independent 3D Schrödinger Equation

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{r}) + U(\mathbf{r})\psi(\mathbf{r}) = E\psi(\mathbf{r})$$

$U(\mathbf{r}) = U(r)$ 일 경우

변수분리

$$\psi(\mathbf{r}) = \psi(r, \theta, \phi) = R(r)\Theta(\theta)\Phi(\phi)$$

$$\nabla^2 = \frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \left(\frac{\partial^2}{\partial \theta^2} + \cot \theta \frac{\partial}{\partial \theta} + \csc^2 \theta \frac{\partial^2}{\partial \phi^2} \right)$$





When Potential Depends on r Only


$$\frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2}$$
$$= -\sin^2 \theta \left\{ \frac{r^2}{R} \left(\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} \right) + \frac{1}{\Theta} \left(\frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} \right) + \frac{2mr^2}{\hbar^2} [E - U(r)] \right\}$$
$$= -m_l^2$$

m_l Magnetic quantum number





When Potential Depends on r Only



$$\frac{d^2\Phi}{d\phi^2} = -m_l^2\Phi(\phi)$$

$$\Phi(\phi) = \frac{1}{\sqrt{2\pi}} e^{im_l\phi}$$

$$m_l = \dots, -2, -1, 0, 1, 2, \dots$$



When Potential Depends on r Only

$$\frac{r^2}{R} \left(\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} \right) + \frac{2mr^2}{\hbar^2} [E - U(r)] = -\frac{1}{\Theta} \left(\frac{d^2 \Theta}{d\theta^2} + \cot \theta \frac{d\Theta}{d\theta} \right) + \frac{m_l^2}{\sin^2 \theta}$$
$$= l(l+1)$$

l Orbital quantum number





When Potential Depends on r Only

$$\frac{d^2\Theta}{d\theta^2} + \cot\theta \frac{d\Theta}{d\theta} - m_l^2 \csc^2\theta \Theta(\theta) = -l(l+1)\Theta(\theta)$$
$$l = 0, 1, 2, \dots$$

$$l \geq |m_l|$$

Associated Legendre polynomials



Table 8.2 Some Associated Legendre Polynomials $P_\ell^{m_\ell}(\cos \theta)$

$$P_0^0 = 1$$

$$P_1^0 = 2 \cos \theta$$

$$P_1^1 = \sin \theta$$

$$P_2^0 = 4(3 \cos^2 \theta - 1)$$

$$P_2^1 = 4 \sin \theta \cos \theta$$

$$P_2^2 = \sin^2 \theta$$

$$P_3^0 = 24(5 \cos^3 \theta - 3 \cos \theta)$$

$$P_3^1 = 6 \sin \theta(5 \cos^2 \theta - 1)$$

$$P_3^2 = 6 \sin^2 \theta \cos \theta$$

$$P_3^3 = \sin^3 \theta$$

© 2005 Brooks/Cole - Thomson





$$\Theta(\theta)\Phi(\phi) = Y_l^{m_l}(\theta, \phi)$$

Spherical harmonics

Table 8.3 The Spherical Harmonics $Y_\ell^{m_\ell}(\theta, \phi)$

$$Y_0^0 = \frac{1}{2\sqrt{\pi}}$$

$$Y_1^0 = \frac{1}{2}\sqrt{\frac{3}{\pi}} \cdot \cos \theta$$

$$Y_1^{\pm 1} = \mp \frac{1}{2}\sqrt{\frac{3}{2\pi}} \cdot \sin \theta \cdot e^{\pm i\phi}$$

$$Y_2^0 = \frac{1}{4}\sqrt{\frac{5}{\pi}} \cdot (3 \cos^2 \theta - 1)$$

$$Y_2^{\pm 1} = \mp \frac{1}{2}\sqrt{\frac{15}{2\pi}} \cdot \sin \theta \cdot \cos \theta \cdot e^{\pm i\phi}$$

$$Y_2^{\pm 2} = \frac{1}{4}\sqrt{\frac{15}{2\pi}} \cdot \sin^2 \theta \cdot e^{\pm 2i\phi}$$

$$Y_3^0 = \frac{1}{4}\sqrt{\frac{7}{\pi}} \cdot (5 \cos^3 \theta - 3 \cos \theta)$$

$$Y_3^{\pm 1} = \mp \frac{1}{8}\sqrt{\frac{21}{\pi}} \cdot \sin \theta \cdot (5 \cos^2 \theta - 1) \cdot e^{\pm i\phi}$$

$$Y_3^{\pm 2} = \frac{1}{4}\sqrt{\frac{105}{2\pi}} \cdot \sin^2 \theta \cdot \cos \theta \cdot e^{\pm 2i\phi}$$

$$Y_3^{\pm 3} = \mp \frac{1}{8}\sqrt{\frac{35}{\pi}} \cdot \sin^3 \theta \cdot e^{\pm 3i\phi}$$

© 2005 Brooks/Cole - Thomson





Hydrogen-like Atom

Radial wave equation

$$-\frac{\hbar^2}{2m} \left(\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} \right) + \frac{l(l+1)\hbar^2}{2mr^2} R(r) + U(r)R(r) = ER(r)$$

For $U(r) = -\frac{Ze^2}{4\pi\epsilon_0 r},$

$$E_n = -\frac{e^2 Z^2}{8\pi\epsilon_0 a_o n^2} = -\frac{me^4 Z^2}{32\pi^2 \epsilon_0^2 \hbar^2 n^2} = -13.6\text{eV} \cdot \frac{Z^2}{n^2}, \quad n = 1, 2, 3, \dots$$

$$\left(a_o = \frac{\hbar^2}{4\pi\epsilon_0 me^2} \quad \text{Bohr radius} \right)$$

Principal quantum number





Table 8.4 The Radial Wavefunctions $R_{n\ell}(r)$ of Hydrogen-like Atoms for $n = 1, 2,$ and 3

n	ℓ	$R_{n\ell}(r)$
1	0	$\left(\frac{Z}{a_0}\right)^{3/2} 2e^{-Zr/a_0}$
2	0	$\left(\frac{Z}{2a_0}\right)^{3/2} \left(2 - \frac{Zr}{a_0}\right) e^{-Zr/2a_0}$
2	1	$\left(\frac{Z}{2a_0}\right)^{3/2} \frac{Zr}{\sqrt{3} a_0} e^{-Zr/2a_0}$
3	0	$\left(\frac{Z}{3a_0}\right)^{3/2} 2 \left[1 - \frac{2Zr}{3a_0} + \frac{2}{27} \left(\frac{Zr}{a_0}\right)^2\right] e^{-Zr/3a_0}$
3	1	$\left(\frac{Z}{3a_0}\right)^{3/2} \frac{4\sqrt{2}}{3} \frac{Zr}{a_0} \left(1 - \frac{Zr}{6a_0}\right) e^{-Zr/3a_0}$
3	2	$\left(\frac{Z}{3a_0}\right)^{3/2} \frac{2\sqrt{2}}{27\sqrt{5}} \left(\frac{Zr}{a_0}\right)^2 e^{-Zr/3a_0}$

© 2005 Brooks/Cole - Thomson





Hydrogen-like Atom

$$E_n = -\frac{me^4 Z^2}{32\pi^2 \epsilon_0^2 \hbar^2 n^2} = -13.6\text{eV} \cdot \frac{Z^2}{n^2},$$

$$\Psi_{nlm_l}(r, \theta, \phi, t) = R_{nl}(r) Y_l^{m_l}(\theta, \phi) e^{-i\frac{E_n}{\hbar}t}$$

$n = 1, 2, 3, \dots$ Principal quantum number

$l = 0, 1, 2, \dots, (n-1)$ Orbital quantum number

$m_l = -l, (-l+1), \dots, -2, -1, 0, 1, 2, \dots, (l-1), l$

$m_s = -\frac{1}{2}, \frac{1}{2}$ Spin magnetic quantum number



표 6.1 수소원자의 규격화된 파동함수; $n = 1, 2, 3$

n	l	m_l	$\Phi(\phi)$	$\Theta(\theta)$	$R(r)$	$\psi(r, \theta, \phi)$
1	0	0	$\frac{1}{\sqrt{2\pi}}$	$\frac{1}{\sqrt{2}}$	$\frac{2}{a_0^{3/2}} e^{-r/a_0}$	$\frac{1}{\sqrt{\pi} a_0^{3/2}} e^{-r/a_0}$
2	0	0	$\frac{1}{\sqrt{2\pi}}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2\sqrt{2} a_0^{3/2}} \left(2 - \frac{r}{a_0}\right) e^{-r/2a_0}$	$\frac{1}{4\sqrt{2\pi} a_0^{3/2}} \left(2 - \frac{r}{a_0}\right) e^{-r/2a_0}$
2	1	0	$\frac{1}{\sqrt{2\pi}}$	$\frac{\sqrt{6}}{2} \cos \theta$	$\frac{1}{2\sqrt{6} a_0^{3/2}} \frac{r}{a_0} e^{-r/2a_0}$	$\frac{1}{4\sqrt{2\pi} a_0^{3/2}} \frac{r}{a_0} e^{-r/2a_0} \cos \theta$
2	1	± 1	$\frac{1}{\sqrt{2\pi}} e^{\pm i\phi}$	$\frac{\sqrt{3}}{2} \sin \theta$	$\frac{1}{2\sqrt{6} a_0^{3/2}} \frac{r}{a_0} e^{-r/2a_0}$	$\frac{1}{8\sqrt{\pi} a_0^{3/2}} \frac{r}{a_0} e^{-r/2a_0} \sin \theta e^{\pm i\phi}$
3	0	0	$\frac{1}{\sqrt{2\pi}}$	$\frac{1}{\sqrt{2}}$	$\frac{2}{81\sqrt{3} a_0^{3/2}} \left(27 - 18 \frac{r}{a_0} + 2 \frac{r^2}{a_0^2}\right) e^{-r/3a_0}$	$\frac{1}{81\sqrt{3\pi} a_0^{3/2}} \left(27 - 18 \frac{r}{a_0} + 2 \frac{r^2}{a_0^2}\right) e^{-r/3a_0}$
3	1	0	$\frac{1}{\sqrt{2\pi}}$	$\frac{\sqrt{6}}{2} \cos \theta$	$\frac{4}{81\sqrt{6} a_0^{3/2}} \left(6 - \frac{r}{a_0}\right) \frac{r}{a_0} e^{-r/3a_0}$	$\frac{\sqrt{2}}{81\sqrt{\pi} a_0^{3/2}} \left(6 - \frac{r}{a_0}\right) \frac{r}{a_0} e^{-r/3a_0} \cos \theta$
3	1	± 1	$\frac{1}{\sqrt{2\pi}} e^{\pm i\phi}$	$\frac{\sqrt{3}}{2} \sin \theta$	$\frac{4}{81\sqrt{6} a_0^{3/2}} \left(6 - \frac{r}{a_0}\right) \frac{r}{a_0} e^{-r/3a_0}$	$\frac{1}{81\sqrt{\pi} a_0^{3/2}} \left(6 - \frac{r}{a_0}\right) \frac{r}{a_0} e^{-r/3a_0} \sin \theta e^{\pm i\phi}$
3	2	0	$\frac{1}{\sqrt{2\pi}}$	$\frac{\sqrt{10}}{4} (3 \cos^2 \theta - 1)$	$\frac{4}{81\sqrt{30} a_0^{3/2}} \frac{r^2}{a_0^2} e^{-r/3a_0}$	$\frac{1}{81\sqrt{6\pi} a_0^{3/2}} \frac{r^2}{a_0^2} e^{-r/3a_0} (3 \cos^2 \theta - 1)$
3	2	± 1	$\frac{1}{\sqrt{2\pi}} e^{\pm i\phi}$	$\frac{\sqrt{15}}{2} \sin \theta \cos \theta$	$\frac{4}{81\sqrt{30} a_0^{3/2}} \frac{r^2}{a_0^2} e^{-r/3a_0}$	$\frac{1}{81\sqrt{\pi} a_0^{3/2}} \frac{r^2}{a_0^2} e^{-r/3a_0} \sin \theta \cos \theta e^{\pm i\phi}$
3	2	± 2	$\frac{1}{\sqrt{2\pi}} e^{\pm 2i\phi}$	$\frac{\sqrt{15}}{4} \sin^2 \theta$	$\frac{4}{81\sqrt{30} a_0^{3/2}} \frac{r^2}{a_0^2} e^{-r/3a_0}$	$\frac{1}{162\sqrt{\pi} a_0^{3/2}} \frac{r^2}{a_0^2} e^{-r/3a_0} \sin^2 \theta e^{\pm 2i\phi}$





Table 8.5 Spectroscopic Notation for Atomic Shells and Subshells

n	Shell Symbol	ℓ	Shell Symbol
1	K	0	s
2	L	1	p
3	M	2	d
4	N	3	f
5	O	4	g
6	P	5	h
...		...	

© 2005 Brooks/Cole - Thomson

Table 8-5, p.280





표 6.2 원자 전자 상태

	$l = 0$	$l = 1$	$l = 2$	$l = 3$	$l = 4$	$l = 5$
$n = 1$	$1s$					
$n = 2$	$2s$	$2p$				
$n = 3$	$3s$	$3p$	$3d$			
$n = 4$	$4s$	$4p$	$4d$	$4f$		
$n = 5$	$5s$	$5p$	$5d$	$5f$	$5g$	
$n = 6$	$6s$	$6p$	$6d$	$6f$	$6g$	$6h$





Angular Momentum



Fingers of right hand in direction of rotational motion

© 2005 Brooks/Cole - Thomson





When Potential Depends on r Only

Radial wave equation

$$-\frac{\hbar^2}{2m} \left(\frac{d^2 R}{dr^2} + \frac{2}{r} \frac{dR}{dr} \right) + \frac{l(l+1)\hbar^2}{2mr^2} R(r) + U(r)R(r) = ER(r)$$

$$K_{orb} = \frac{1}{2} mv^2 = \frac{L^2}{2mr^2}$$

$$L = \sqrt{l(l+1)}\hbar$$





Angular Momentum

$$\hat{\mathbf{L}} = \hat{\mathbf{r}} \times \hat{\mathbf{p}}$$

$$\hat{\mathbf{L}} = \begin{vmatrix} \mathbf{a}_x & \mathbf{a}_y & \mathbf{a}_z \\ x & y & z \\ -i\hbar\partial/\partial x & -i\hbar\partial/\partial y & -i\hbar\partial/\partial z \end{vmatrix}$$

$$\hat{L}_x = i\hbar \left(\sin\phi \frac{\partial}{\partial\theta} + \cot\theta \cos\phi \frac{\partial}{\partial\phi} \right)$$

$$\hat{L}_y = -i\hbar \left(\cos\phi \frac{\partial}{\partial\theta} - \cot\theta \sin\phi \frac{\partial}{\partial\phi} \right)$$

$$\hat{L}_z = -i\hbar \frac{\partial}{\partial\phi}$$





Angular Momentum

$$\hat{L}^2 Y_l^{m_l}(\theta, \phi) = l(l+1)\hbar^2 Y_l^{m_l}(\theta, \phi)$$

$$\hat{L}_z Y_l^{m_l}(\theta, \phi) = m_l \hbar Y_l^{m_l}(\theta, \phi)$$

$$L = \sqrt{l(l+1)}\hbar$$

$$L_z = m_l \hbar$$





Space Quantization




그림 6.4 궤도 각운동량의 공간 양자화






그림 6.6 각운동량 벡터 L 은 z 축 주위로 계속하여 세차 운동을 한다.





(a)

그림 6.5 불확정성 원리에 의해 각운동량 벡터 L 은 공간에서 확정된 방향을 가질 수 없다.



(b)






그림 6.8 다양한 양자 상태에서의 수소원자 지름 파동함수의 핵으로부터의 거리에 따른 변화.




그림 6.12








그림 6.12 계속
Seoul National University






$$n = 2, \ell = 1, m_\ell = \pm 1$$

(a)



$$n = 3, \ell = 1, m_\ell = 0$$

(b)



$$n = 3, \ell = 2, m_\ell = 0$$

(c)

© 2005 Brooks/Cole - Thomson







그림 6.9 구면 극좌표계에서의 부피 요소 dV



$$P(r)dr = |\psi|^2 4\pi r^2 dr$$

$$P(r) = r^2 |R(r)|^2$$



핵으로부터의 거리가 r 과 $r+dr$ 사이인 공 껍질에서 수소원자의 전자를 발견할 확률은 $P(r)dr$ 이다.

© 2005 Brooks/Cole - Thomson








그림 6.11 그림 6.8에서 나타낸 상태들에 대한, 핵으로부터의 거리 r 과 $r+dr$ 사이에서 수소원자의 전자를 발견할 확률.



(a)



(b)

© 2005 Brooks/Cole - Thomson

