# Stoichiometry of Biochemical Reactions I

# Today's lecture

- Biochemical reaction stoichiometry
- Cell yield
- Half reactions

#### Oxidation-reduction (redox) reactions

- Involves changes in the oxidation state
- If there are oxidizing species, there should be reducing species as well
  - Total # of electrons are conserved
  - Oxidizing species e<sup>-</sup> donor / Reducing species –
     e<sup>-</sup> acceptor
- Essential for life many important biological processes involve redox reactions
  - Involves large free energy change ( $\Delta G$ )

#### Biological benefit of redox reactions

#### **Examples:**

photosynthesis of glucose:

$$6CO_2 + 6H_2O + sunlight \rightarrow C_6H_{12}O_6 + 6O_2$$

 $\Delta G^0 = +2880 \text{ kJ/mol } C_6 H_{12} O_6$ 

Sunlight energy is converted to chemical energy (storable)

respiration using glucose:

$$C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$$

 $\Delta G^0 = -2880 \text{ kJ/mol C}_6 H_{12} O_6$ 

Chemical energy is released to be used to do something

### Stoichiometry

- "An aspect of chemistry concerned with mole relationships among reactants and products"
- Simply put, balancing chemical reactions
- Based on mass conservation
  - Conservation of elements
  - Conservation of electrons (for redox reactions)

#### What can we do with stoichiometry?

#### A balanced biochemical reaction example

(ethanol fermentation of glucose at  $f_s$ =0.22)

$$0.0417C_6H_{12}O_6 + 0.011NH_4^+ + 0.011HCO_3^-$$

$$\rightarrow$$
 0.011C<sub>5</sub>H<sub>7</sub>O<sub>2</sub>N + 0.065C<sub>2</sub>H<sub>5</sub>OH + 0.076CO<sub>2</sub> + 0.044H<sub>2</sub>O [Bacterial cells]

#### Available information:

Using 1 g (or 1 mole) of glucose,

- How much ethanol can be produced?
- How much nutrients (NH<sub>4</sub>-N) are required?
- How much biomass is produced?
- How much alkalinity is consumed?

#### Cell formula


- Most common: C<sub>5</sub>H<sub>7</sub>O<sub>2</sub>N
- What is the theoretical COD-per-weight value for a bacterial cell (i.e., g COD / g cells)?
  - Once you get trained with the stoichiometry, you will be able to write the balanced reaction as:

$$C_5H_7O_2N + 5O_2 \rightarrow 5CO_2 + NH_3 + 2H_2O$$

Then you get: 1.42 g COD/g cells

You will find this value useful in the future!

# Substrate partitioning



## Cell yield

True yield, Y
 Y = (g cells produced) / (g substrate utilized)

• Conversion of  $f_s^0$  to Y:

$$Y = f_s^{\ 0} \frac{(M_c \ g \ cells/mole \ cells)}{(n_e \ e^- \ eq \ cells/mole \ cells)(8 \ g \ COD/e^- \ eq \ donor)}$$

For 
$$C_5H_7O_2N$$
,  $M_c = 113$  g/mole;  $n_e = 20 e^-$  eq/mole (see Table 2.3) then:  $Y(in \ g \ cells/g \ COD) = 0.706 f_s^{\ 0}$ 

#### Microbial growth rate

$$\frac{dX_a}{dt} = Y \left(\frac{-dS}{dt}\right) - bX_a$$
growth decay

 $X_q$  = active biomass concentration [M/L<sup>3</sup>]

S = substrate concentration [M/L<sup>3</sup>]

Y = true yield [M/M]

b = decay rate [1/T]

#### **Decay**

- Generally assumed to be proportional to the amount of cells
- A gross factor for anything that leads to decrease in cell biomass
- Decay = endogenous respiration (+ predation)
  - Endogenous respiration: use of cell matter for maintenance of cell functions (motility, repair, resynthesis, osmotic regulation, transport, compensate heat loss, ...)


## Net yield

• Net yield,  $Y_n$  $Y_n = (g net cell growth) / (g substrate utilized)$ 

$$= \frac{dX_a / dt}{-dS / dt}$$

$$= Y - b \frac{X_a}{-dS / dt}$$

# Net yield



Log (exponential) growth:

Stationary phase:

Death phase:

Bacterial growth curve for pure culture

### **Net yield**

• Electron partitioning considering net yield,  $Y_n$ :

$$f_s^0 \rightarrow f_s$$
  $(f_s < f_s^0)$   
 $f_e^0 \rightarrow f_e$   $(f_e > f_e^0)$ 

still, 
$$f_s + f_e = 1$$

#### **Energy reactions**

Microorganisms need energy for growth and maintenance

 $\Delta G^0$  (in kJ/mole glucose)

Aerobic oxidation:

$$C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$$
 -2,880

Denitrification:

$$5C_6H_{12}O_6 + 24NO_3^- 24H^+ \rightarrow 30CO_2 + 42H_2O + 12N_2$$
 -2,720

Sulfate reduction:

$$2C_6H_{12}O_6 + 6SO_4^{2-} \rightarrow 12CO_2 + 12H_2O + 3H_2S + 3HS^{-}$$
 -492

Methanogenesis:

$$C_6H_{12}O_6 \rightarrow 3CO_2 + 3H_2O$$
 -428  
 $(C_6H_{12}O_6 + 3CO_2 \rightarrow 6CO_2 + 3H_2O)$ 

**Ethanol fermentation:** 

$$C_6H_{12}O_6 \rightarrow 2CO_2 + 2C_2H_5OH$$
 -244 15

#### Half reactions

- For complex biochemical redox reactions, it is easier to use half reaction approach
- The oxidation reaction for an electron donor and the reduction reaction for an electron acceptor can be splitted
- Usually written as a reduction reaction (see [Table 2.2] & [Table 2.3])

#### Half reactions

- **Step 1** Write oxidized form on the left and reduced form on the right
- **Step 2** Add other species involved in the reaction
- **Step 3** Balance the reaction for all elements except for oxygen and hydrogen
- Step 4 Balance oxygen using water
- Step 5 Balance hydrogen using H<sup>+</sup>
- Step 6 Balance charge using e
- **Step 7** Convert the equation to the e-equivalent form