

Toxic and recalcitrant compound removal

Toxic & recalcitrant organic compounds


- **Xenobiotic compounds: man-made**
 - Many are resistant to biodegradation and have potential toxicity to ecosystem & human health
- **Refractory, recalcitrant – not easily biodegradable**
- Some specific microorganisms may have the ability to degrade toxic and recalcitrant compounds

Examples of toxic & recalcitrant organics

| Type of waste | Types of organic compounds |
|--------------------------------------|--|
| Petroleum | alkanes; alkenes; polyaromatic hydrocarbons; monocyclic aromatics – benzene, toluene, ethylbenzene, xylenes; naphthenes |
| Non-halogenated solvents | alcohols; ketones; esters; ethers; aromatic and aliphatic hydrocarbons; glycols; amines |
| Halogenated solvents | chlorinated methanes – methylene chloride, chloroform, carbon tetrachloride; chlorinated ethenes – tetrachloroethene, trichloroethene; chlorinated ethanes – trichloroethane; chlorinated benzenes |
| Insecticides, herbicides, fungicides | organochloride compounds; organophosphate compounds; carbamate esters; phenyl ethers; creosotes; chlorinated phenols |
| Munitions and explosives | nitroaromatics – trinitrotoluene; nitramines; nitrate esters |
| Industrial intermediates | phthalate esters; benzene; phenol; chlorobenzenes; chlorophenols; xylenes |
| Transformer and hydraulic fluids | polychlorinated biphenyls |
| Production byproducts | dioxin, furans |

Abiotic losses in WWTP

- May be significant for many toxic & recalcitrant compounds
- Adsorption to biomass in secondary treatment
 - Removed from wastewater as sludge
 - Issues with sludge application and disposal
- Volatilization: released to the atmosphere



Fate of non-biodegradable and non-volatile organic compounds during conventional wastewater treatment


Heidler & Halden (2008) *ES&T* 42:6324-6332.

Biodegradation pathways (1)

- **Compound serving as a growth substrate**
 - Complete mineralization or transformation to a different compound (hopefully less or non-toxic)
 - Aerobic degradation usually more significant than anaerobic
 - Aerobic degradation works for many non-halogenated organic compounds, but not often for halogenated compounds

Aerobic biodegradation of aromatics

- Aromatic rings are relatively stable
 - Ring cleavage is often the major challenge
- Some specific enzymes are able to insert oxygen into the aromatic ring to initiate the degradation



Proposed benzene degradation pathway by Bacillus pumilus MVSV3
(Surendra et al., 2017, Braz. Arch. Biol. Technol.)


Biodegradation pathways (2)

- **Compound as an e⁻ acceptor**

- [reductive dehalogenation]

- Under sulfidogenic/methanogenic condition (highly reduced condition)
- Uses H₂ as an e⁻ donor – substitution of a halogen (Cl, F, ..) with H in the organic molecule
- Reductive dechlorination extensively studied
 - PCE, TCE, PCP, PCBs, etc.

PCE dechlorination pathway




- Ethene can be easily mineralized by other microorganisms
- Dechlorination more challenging for DCE/VC than PCE/TCE
 - Only a very limited number of species can completely reduce PCE to ethene
 - VC is more toxic than PCE/TCE – possibility of increased toxicity by biotransformation
 - So should confirm that the species capable of complete PCE/TCE reduction to ethene are present at the site for bioremediation

Biodegradation pathways (3)

- **Cometabolism**


- Transformation of a compound by a microorganism that is unable to use the compound as a carbon or energy source
- No microorganism growth by cometabolism
- Degradation pathway for chlorinated organic compounds under aerobic condition
- By bacteria producing nonspecific mono-oxygenase or dioxygenase enzymes
- Example organisms: methanotrophic bacteria, phenol/toluene oxidizers

Methanotroph transforming TCE by cometabolism



- Methane is a growth substrate
- MMO catalyzes methane oxidation to methanol
- The MMO produced is also capable of catalyzing TCE oxidation to TCE-epoxide
- In pure culture, TCE-epoxide accumulates, but in the environment, it may be further degraded by other microorganisms to be mineralized

Toluene injection for treatment of TCE in GW



Kuo et al., 2004, Water Res.

- Toluene & O₂ injection to promote growth of toluene-oxidizing bacteria
- TCE biodegradation by cometabolism
- >90% TCE removal observed
- Constraint of adding a pollutant to remove another

Fig. 1. Schematic diagrams of Pilot A and venturi tube used for toluene-vapor injection.