

Water constituents

What's in water

- **Microorganisms**

- Native vs. introduced; Oligotrophic vs. eutrophic
- Millions to billions per mL or g
- Complex consortia
 - Species makeup: f(T, O₂, pH, nutrients, etc.)
- Indicators & pathogens

- **Dissolved inorganics**

- Major vs. trace
- Nutrient vs. contaminant

What's in water

- **Dissolved organics (DOM)**
 - Simple vs. polymeric
 - Biodegradable vs. recalcitrant
 - Natural vs. anthropogenic
- **Suspended matter**
 - Operationally define: retained on 0.45 μm -filter
 - Mineral colloids; organic polymers; bacteria
 - Aggregation & sorption
 - Contaminant sink, transport consequences

Microbes

- **$C_5H_7O_2N$ – approx. microbial chemical formula**
 - Dry weight basis; Cell is composed of ~90% water by weight
- **Other major constituents**
 - P: ~0.07 mole/mole cell material
 - S, Mg, Ca, K, Na: ~0.02-0.03 mole/mole cell material
 - Fe: more than trace, less than major
- **Trace constituents**
 - Cr, Co, Mn, Mo, Ni, Se, W, V, Zn
 - Constituents of enzyme systems
 - Need a little, but *too much is toxic*

Dissolved inorganics

- **Major constituents of surface & groundwaters (ppm+)**
 - Na^+ , K^+ , Ca^{2+} , Mg^{2+} , Cl^- , HCO_3^- , SO_4^{2-} , SiO_2
- **Others important to life (ppb~ppm)**
 - O_2 ; Fe; N (NO_3^- , NH_3); P (PO_4^{3-})
 - N & P: rapid uptake by biota
 - P: strongly sorbed to minerals
 - Fe: low solubility in aerobic waters (Fe^{3+}) and moderate pH
- **Special cases**
 - Oxygen absent: high Fe^{2+} , S^{2-}
 - Low pH – high Fe^{3+}
- **Total dissolved solids (TDS): primarily ions**
 - Surface water < groundwater < seawater

Dissolved inorg. - electroneutrality

- **Electroneutrality principle**

$$\sum \text{cations (in eq/L)} = \sum \text{anions (in eq/L)}$$

- **May use the following condition to determine the accuracy of water ion content analysis**

$$\left| \sum \text{anions} - \sum \text{cations} \right| \leq \left(0.1065 + 0.0155 \sum \text{anions} \right)$$

Dissolved inorg. - electroneutrality

- **Ionic strength**

$$I = \frac{1}{2} \sum_i (C_i \times z_i^2)$$

C_i = concentration of ionic species i (M)
z_i = charge of ionic species i

- Significance: in dilute solutions ($I \sim < 10^{-3}$ M) the ions behave independently of each other, but as ion concentration increases, ion interactions become significant, decreasing the activity of the ions

- **Activity vs. molarity ({ } vs. [])**

$$\{i\} = \gamma_i [i]$$

- $\gamma_i \cong 1$ in dilute solutions (for most natural waters except for seawater, this would be acceptable for crude calculations)
- Güntelberg equation (for $I < 0.1$):

$$\log \gamma_i = - \frac{0.5 z_i^2 I^{0.5}}{1 + I^{0.5}}$$

Organics in natural waters


- **Simple sugars, amino acids, etc.**
 - Concentrations typically very low – easily degraded, assimilated
- **Microbial polymers**
 - Important components of biofilms, flocs, aggregates
 - In some cases may have significant dissolved concentrations
- **Humic substances**
 - Typically the primary component of dissolved & particulate organic matter (DOM, POM)
 - Resistant to degradation
 - Many ions & hydrophobic compounds sorb strongly to humics

Organics in natural waters

- **Anthropogenic organics**
 - >100,000 synthetic chemicals in daily use
 - Pesticides, solvents, dyes, personal care products, anti-fouling agents, additives
 - >300,000,000 tons produced annually
 - Wide varying properties
 - Size, aqueous solubility, volatility, degradability, toxicity



Dissolved organics – humic substances

- **Acidic polymers**
 - ~3-15 mmoles/g, primarily
 - carboxylic
 - Phenolic OH
- **Wide varying composition**
 - There is no “humic molecule”
 - MWs from ~500 to >100,000
- **Molecular volume a function of pH, ionic strength**
 - Compact at high salinity, low pH
 - Extend at low salinity, high pH



Dissolved organics – humic substances

- **Hydrophilic/hydrophobic regions**
 - High affinity for many toxic organics
- **Coat minerals**
- **Photoactive**
- **High metal affinity**



Suspended matter

- **Not dissolved; potentially settleable (usually)**
 - Traditionally considered to be material that retained on a 0.45 μm -filter
 - Colloids may be as small as a few nanometers
- **Includes: mineral colloids, microbes and their debris, organic polymers**
 - Often found as aggregates, flocs that are mixtures of minerals, polymers, and microbial matter
 - Aggregates $>0.45 \mu\text{m}$ may consist of many individual components $\ll 0.45 \mu\text{m}$

Suspended matter

- **Suspended mater influences:**
 - Contaminant transport, scavenging
 - Light attenuation
 - Disinfection efficacy
 - Where the material settles

Aqueous chemistry parameters

- **Units**

- Mass/vol kg/m³, mg/L, ...
- #/vol mole/m³, # of organisms/mL
- Transferable electrons or protons/vol equiv/m³, equiv/L
- Mole fraction mole/Σmole

- **Why different units?**

- Engineers vs. chemists
- Specific needs of the problem

- **Aggregate parameters**

- BOD, COD, TOC
- Hardness
- Total PCBs

Aqueous chemistry parameters






- **Aggregate parameters**
 - Characterize important properties of mixtures
 - OD (oxygen demand)
 - TO (total organic carbons (C), halides (X))
 - Conduct one analysis instead of many
- **per as**
 - Prevent confusion
 - mg/L as CaCO₃
 - mg/L as N
 - % as P₂O₅ or K₂O

Measures of (oxidizable) organic matter

- **BOD – Biochemical Oxygen Demand**
 - Measure of a water's biologically oxidizable constituents
 - Analyze [DO] in a water sample before & after controlled incubation
 - 5 day incubation is common
- **COD – Chemical Oxygen Demand**
 - Measure of a water's chemically oxidizable constituents
 - 2-3 hour reaction time
 - Generates liquid hazardous wastes
 - Doesn't oxidize organic N
- **TOC, DOC – Total/Dissolved Organic Carbon**
 - Measure of a water's organic carbon content
 - Analyze mass/concentration of CO₂ produced after chemical oxidation of a sample
 - Sampling time – a few minutes

BOD₅, COD & TOC

Compound	Formula	MW	BOD ₅	COD	TOC	COD/TOC	TOC/MW	COD/MW
Methane	CH ₄	16	??	64	12	5.3	0.75	4.0
MTBE	C ₅ H ₁₂ O	88	~0	240	60	4.0	0.68	2.7
Benzene	C ₆ H ₆	78	??	240	72	3.3	0.92	3.1
Glucose	C ₆ H ₁₂ O ₆	192	~192	192	72	2.7	0.38	1.0



[MTBE]

Aqueous chemistry parameters – pH

- **$\text{pH} = -\log_{10}\{\text{H}^+\} \approx -\log_{10}[\text{H}^+]$**
 - For most natural waters $5 < \text{pH} < 9$
 - Most aquatic life adapted for this range
 - $K_w = \{\text{H}^+\}\{\text{OH}^-\} = 10^{-14}$ @25°C
 - $\text{p}K_w = \text{pH} + \text{pOH} \approx 14$
 - Chemical speciation can be highly pH dependent



- If $\text{p}K_a > \text{pH}$, deprotonated (dissociated)
- If $\text{p}K_a < \text{pH}$, protonated (associated)