

Introduction to Electromagnetism

Time-Varying Fields and Maxwell's Equations

(7-6, 7-7)

Yoonchan Jeong

School of Electrical Engineering, Seoul National University

Tel: +82 (0)2 880 1623, Fax: +82 (0)2 873 9953

Email: yunchan@snu.ac.kr

Farewell to Electrostatic Model

Electrostatic model

$$\nabla \times \mathbf{E} = 0$$

$$\nabla \cdot \mathbf{D} = \rho$$

$$\mathbf{D} = \varepsilon \mathbf{E}$$

Magnetostatic model

$$\nabla \times \mathbf{H} = \mathbf{J}$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\mathbf{B} = \mu \mathbf{H}$$

Maxwell's equations

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad \textit{Faraday's law}$$

$$\nabla \times \mathbf{H} = \mathbf{J} + \frac{\partial \mathbf{D}}{\partial t} \quad \textit{Ampère's law}$$

$$\nabla \cdot \mathbf{D} = \rho \quad \textit{Gauss's law}$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\mathbf{D} = \varepsilon \mathbf{E} = \varepsilon_0 \mathbf{E} + \mathbf{P} \quad \textit{Constitutive relations}$$

$$\mathbf{H} = \frac{1}{\mu} \mathbf{B} = \frac{1}{\mu_0} \mathbf{B} - \mathbf{M}$$

Source-Free Wave Equations

Source free, simple medium: $\rho = 0$ & $\mathbf{J} = 0$ ($\sigma = 0$)

→ *Linear, isotropic and homogeneous*

Maxwell's equations

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

$$\nabla \times \mathbf{H} = \mathbf{J} + \frac{\partial \mathbf{D}}{\partial t}$$

$$\nabla \cdot \mathbf{D} = \rho$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\mathbf{D} = \epsilon \mathbf{E} = \epsilon_0 \mathbf{E} + \mathbf{P}$$

$$\mathbf{H} = \frac{1}{\mu} \mathbf{B} = \frac{1}{\mu_0} \mathbf{B} - \mathbf{M}$$



$$\nabla \times \mathbf{E} = -\mu \frac{\partial \mathbf{H}}{\partial t}$$

$$\nabla \times \mathbf{H} = \epsilon \frac{\partial \mathbf{E}}{\partial t}$$

$$\nabla \cdot \mathbf{E} = 0$$

$$\nabla \cdot \mathbf{H} = 0$$

$$\nabla \times \nabla \times \mathbf{E} = -\mu \frac{\partial}{\partial t} (\nabla \times \mathbf{H}) = -\mu \epsilon \frac{\partial^2 \mathbf{E}}{\partial t^2}$$

$$= \nabla(\nabla \cdot \mathbf{E}) - \nabla^2 \mathbf{E}$$

$$\rightarrow \nabla^2 \mathbf{E} - \mu \epsilon \frac{\partial^2 \mathbf{E}}{\partial t^2} = 0$$

$$u = 1 / \sqrt{\mu \epsilon}$$

→ *Phase velocity*

$$\rightarrow \nabla^2 \mathbf{E} - \frac{1}{u^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} = 0$$

$$\rightarrow \nabla^2 \mathbf{H} - \frac{1}{u^2} \frac{\partial^2 \mathbf{H}}{\partial t^2} = 0$$

→ *Homogeneous wave equations* 3

Electromagnetic Waves

Wave equations:

$$\nabla^2 \mathbf{E} - \mu\epsilon \frac{\partial^2 \mathbf{E}}{\partial t^2} = 0, \quad \nabla^2 \mathbf{H} - \mu\epsilon \frac{\partial^2 \mathbf{H}}{\partial t^2} = 0 \quad (\text{Homogeneous and no source})$$

Plane waves:


$$\psi = e^{i(\omega t - \mathbf{k} \cdot \mathbf{r})}, \quad |\mathbf{k}| = \omega \sqrt{\mu\epsilon}$$

$$\text{e.g. } f(x,t) = f(x - \delta x, t - \delta t)$$

Phase velocity:


$$\omega t - \mathbf{k} \cdot \mathbf{r} = \text{const}, \quad u_p = \frac{\omega}{k} = \frac{1}{\sqrt{\mu\epsilon}},$$

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}} = 2.997930 \times 10^8 \quad (\text{m/s})$$




Doppler Effect in EM Waves

Stationary source:



$$v = 0$$

Moving source:



Recall: Speed of light

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}} = 2.997930 \times 10^8 \text{ m/s} \quad \leftarrow \text{Is this frequency dependent?}$$

\rightarrow *Relative to ether?*


Incoming wave (blue-shifted): $c'_{in} = c + v$

Outgoing wave (red shifted): $c'_{out} = c - v$

} Is this true?

Special & general relativity introduced by A. Einstein!!

Electromagnetic Spectrum



Visible light (400 nm ~ 700 nm)

