


CHAPTER 1. FIRST-ORDER ODE

2017.3
서울대학교
조선해양공학과


노명일

※ 본 강의 자료는 이규열, 장범선 교수님께서 만드신 자료를 바탕으로 일부 편집한 것입니다.


Why Mathematics?



How long does it take to cool down the hot water



Increase of population in the future



Changing **rate** of water temperature \propto Temperature difference between water and outside

$$\frac{dT(t)}{dt} = k(T - T_A), k < 0$$

T : Water temperature
 T_A : Outside temperature (constant)

<Newton's law of cooling>

Increasing **rate** of population \propto Present Population

$$\frac{dy(t)}{dt} = k \cdot y(t)$$

y : population
 t : time
 k : proportional constant

<Malthus's population dynamics>

Why do we need to study "Engineering Mathematics"?



Real World



Explain or Predict


Solution

$$z(t) = e^{-\alpha t} (A \cos \omega t + B \sin \omega t) + C \cos \omega_0 t$$



Idealization or Simplification*

- Insight
- Constitutive Relations




Linearization

- Taylor Series




Mathematical Model

$$mz'' + cz' + kz = F_0 \cos \omega_0 t$$



Problem-Solving Abilities



- Linearly Dependent / Independent
- Basis
- Orthogonality
- Linear Combination

* Keener, J.P., Principles of Applied Mathematics, Westview Press, 2000, p.xi : ...there is the goal to explain or predict the behavior of some physical situation. One begins by constructing a mathematical model which captures the essential features of the problem without asking its content with overwhelming detail

1.1 Basic Concepts. Modeling

Modeling

❖ The typical steps of modeling in detail

Step 1. The transition from the physical situation to its mathematical formulation

Step 2. The solution by a mathematical method

Step 3. The physical interpretation of differential equations and their applications

1.1 Basic Concepts. Modeling

- ❖ **Differential Equation (미분방정식):** An equation containing derivatives of an unknown function



- ❖ **Ordinary Differential Equation:** An equation that contains one or several derivatives of an unknown function (y) of **one independent variable (x)**

ex) $y' = \cos x$, $y'' + 9y = e^{-2x}$, $y'y''' - \frac{3}{2}(y')^2 = 0$

- ❖ **Partial Differential Equation:** An equation involving partial derivatives of an unknown function (u) of **two or more variables (x, y)**

ex) $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$

1.1 Basic Concepts. Modeling

❖ **Order (계):** The highest derivative of the unknown function

ex) (1) $y' = \cos x \Rightarrow$ First order

(2) $y'' + 9y = e^{-2x} \Rightarrow$ Second order

(3) $y' y''' - \frac{3}{2}(y')^2 = 0 \Rightarrow$ Third order

❖ **First-order ODE:** Equations contain only the first derivative y' and may contain y and any given functions of x

- Explicit (양함수) Form: $y' = f(x, y)$
- Implicit (음함수) Form: $F(x, y, y') = 0$

1.1 Basic Concepts. Modeling

- ❖ **Solution:** Functions that make the equation hold true
 - **General Solution (일반해)**
: a solution containing an **arbitrary constant**
 - **Particular Solution (특수해)**
: a solution that we choose a specific constant
 - **Singular Solution (Problem 16) (특이해)**
: an additional solution that cannot be obtained from the general solution
 - **Ex. (Problem 16) ODE** : $(y')^2 - xy' + y = 0$
 - General solution** : $y = cx - c^2$
 - Particular solution** : $y = 2x - 4$
 - Singular solution** : $y = x^2 / 4$

1.1 Basic Concepts. Modeling

- ❖ **Initial Value Problems (초기값 문제):** An ordinary differential equation together with specified value of the unknown function at a given point in the domain of the solution

$$y' = f(x, y), \quad y(x_0) = y_0$$

- ☑ Ex.4 Solve the initial value problem

$$y' = \frac{dy}{dx} = 3y, \quad y(0) = 5.7$$

Step 1 Find the general solution.

General solution: $y(x) = ce^{3x}$

Step 2 Apply the initial condition. $y(0) = ce^0 = c = 5.7$

Particular solution: $y(x) = 5.7e^{3x}$

1.1 Basic Concepts. Modeling

- ☑ **Ex. 5** Given an amount of a **radioactive substance**, say 0.5 g (gram), **find the amount present at any later time.**

Physical Information.

Experiments show that at each instant a radioactive substance decomposes at a rate proportional to the amount present.

Step 1 Setting up a mathematical model (a differential equation) of the physical process.

By the physical law : $\frac{dy}{dt} \propto -y \quad \Rightarrow \quad \frac{dy}{dt} = -ky$

The initial condition : $y(0) = 0.5$

Step 2 Mathematical solution.

General solution: $y(t) = ce^{-kt}$

Particular solution: $y(0) = ce^0 = c = 0.5 \quad \Rightarrow \quad y(t) = 0.5e^{-kt}$

Always check your result: $\frac{dy}{dt} = -0.5ke^{-kt} = -ky, \quad y(0) = 0.5e^0 = 0.5$

Step 3 Interpretation of result. The limit of y as $t \rightarrow \infty$ is zero.

1.2 Geometric Meaning of $y' = f(x, y)$. Direction Fields, Euler's Method

1.2 Geometric Meaning of $y'=f(x, y)$. Direction Fields, Euler's Method

❖ A first-order ODE $y' = f(x, y)$

: A solution curve (해 곡선) that passes through a point (x_0, y_0) must have, at that point, the slope y' (x_0) equal to the value of f at that point


$$y'(x_0) = f(x_0, y_0)$$

❖ Direction Field (방향장)

- The graph includes pairs of grid points and line segments
- The line segment at grid point coincides with the tangent line to the solution.

❖ Reason of importance of the direction field

- You need not solve a ODE.
- The method shows the whole family of solutions and their typical properties.



Direction field of with three approximate solution $y' = y + x$, curves passing through $(0, 1)$, $(0, 0)$, $(0, -1)$, respectively

1.2 Geometric Meaning of $y'=f(x, y)$. Direction Fields, Euler's Method

❖ Numeric Method by Euler

: yields approximate solution values at equidistant x -values with an initial value x_0 .

$$y(x_0) = y_0$$


$$x_1 = x_0 + h, \quad y_1 = y_0 + hf(x_0, y_0)$$

$$x_2 = x_0 + 2h, \quad y_2 = y_1 + hf(x_1, y_1)$$

$$x_3 = x_0 + 3h, \quad y_3 = y_2 + hf(x_2, y_2)$$

\vdots \vdots

where the step h : a smaller value for greater accuracy e.g. 0.1 or 0.2



First Euler step, showing a solution curve, its tangent at (x_0, y_0) , step h and increment $hf(x_0, y_0)$ in the formula for y_1

1.2 Geometric Meaning of $y'=f(x, y)$. Direction Fields, Euler's Method

- ODE


$$y' = y+x, \quad x=0, \quad y(0)=0, \quad h=0.2$$

- Exact Solution

$$y = e^x + x + 1$$

Euler method for $y'=y+x, y(0)=0$ for $x=0, \dots, 1.0$ with step $h=0.2$

n	x_n	y_n	$y(x_n)$	Error
0	0.0	0.000	0.000	0.000
1	0.2	0.000	0.021	0.021
2	0.4	0.04	0.092	0.052
3	0.6	0.128	0.222	0.094
4	0.8	0.274	0.426	0.152
5	1.0	0.488	0.718	0.230



$$y' = f(x, y) = y+x$$

$$x_1 = x_0 + h, \quad y_1 = y_0 + hf(x_0, y_0) = y_0 + h(y_0 + x_0)$$

$$x_1 = 0 + 0.2 = 0.2$$

$$y_1 = 0 + 0.2 \cdot 0 = 0$$

$$x_2 = x_1 + h, \quad y_2 = y_1 + hf(x_1, y_1) = y_1 + h(y_1 + x_1)$$

$$x_2 = 0.2 + 0.2 = 0.4$$

$$y_2 = 0 + 0.2 \cdot (0 + 0.2) = 0.04$$

$$x_3 = x_2 + h, \quad y_3 = y_2 + hf(x_2, y_2) = y_2 + h(y_2 + x_2)$$

$$x_3 = 0.4 + 0.2 = 0.6$$

$$y_3 = 0.04 + 0.2 \cdot (0.04 + 0.4) = 0.128$$

$$x_4 = x_3 + h, \quad y_4 = y_3 + hf(x_3, y_3) = y_3 + h(y_3 + x_3)$$

$$x_4 = 0.6 + 0.2 = 0.8$$

$$y_4 = 0.128 + 0.2 \cdot (0.128 + 0.6) = 0.274$$

Fig. 9. Euler method: Approximate values in Table 1.1 and solution curve

1.3 Separable ODEs. Modeling

1.3 Separable ODEs. Modeling

❖ Separable Equation (변수분리형 방정식):

A differential equation to be separable **all the y 's in the differential equation must be multiplied by the derivative** and **all the x 's in the differential equation must be on the other side of the equal sign.**

$$g(y)y' = f(x) \quad \Rightarrow \quad g(y)dy = f(x)dx \quad \left(\because y' = \frac{dy}{dx} \right)$$

❖ Method of Separating Variables (변수분리법)

$$g(y)y' = f(x) \quad \Rightarrow \quad \int g(y)dy = \int f(x)dx + c \quad \left(\because \frac{dy}{dx} dx = dy \right)$$

☑ **Ex. 1 Solve** $y' = 1 + y^2$ _____ ●

$$\begin{aligned} \frac{y'}{1+y^2} = 1 & \Rightarrow \frac{dy/dx}{1+y^2} = 1 & \Rightarrow \frac{dy}{1+y^2} = dx \\ & \Rightarrow \int \frac{1}{1+y^2} dy = \int dx + c & \Rightarrow \arctan y = x + c & \Rightarrow y = \tan(x + c) \end{aligned}$$

1.3 Separable ODEs. Modeling

✓ Example

Q?

Solve the IVP (Initial Value Problems).

$$\frac{dy}{dx} = -\frac{x}{y}, \quad y(4) = -3$$

1.3 Separable ODEs. Modeling


☑ Ex. 5: Find the amount of salt in the tank at any time t .

→ the amount of salt in the tank = $y(t)$ in 1,000 gal

- The tank contains 1,000 gal of water in which initially 100 lb of salt is dissolved. → Initial condition $y(0) = 100$ lb
- Brine (소금물) runs in at a rate of 10 gal/min, and each gallon contains 5 lb of dissolved salt. → $y_{\text{inflow}} = 10 \text{ gal/min} \times 5 \text{ lb/gal} = 50 \text{ lb/min}$
- The mixture in the tank is kept uniform by stirring (휘저음).
- Brine runs out at 10 gal/min
→ $y_{\text{outflow}} = 10 \text{ gal/min} \times y/1000 \text{ (lb/gal)} = (y/100) \text{ lb/min}$
- Find the amount of salt in the tank at any time t . —————●

the amount of salt = y lb in 1,000 gal

$$10 \text{ gal/min} \times 5 \text{ lb/gal} = 50 \text{ lb/min}$$



$$10 \text{ gal/min} \times y/1000 \text{ lb/gal} = y/100 \text{ lb/min}$$

1 gal = 3.78 l
1 lb = 0.454 kg

Tank

1.3 Separable ODEs. Modeling

☑ Ex. 5

$$10 \text{ gal/min} \times 5 \text{ lb/gal} \\ = 50 \text{ lb/min}$$

the amount of salt = y lb in 1000 gal



$$10 \text{ gal/min} \times y/1000 \text{ lb/gal} \\ = y/100 \text{ lb/min}$$

Tank

Step 1 Setting up a model.

► Salt's time rate of change = Salt inflow rate – Salt outflow rate **“Balance law”**

$$\text{Salt inflow rate} = 10 \text{ gal/min} \times 5 \text{ lb/gal} = 50 \text{ lb/min}$$

$$\text{Salt outflow rate} = 10 \text{ gal/min} \times y/1000 \text{ lb/gal} = y/100 \text{ lb/min}$$

$$(dy/dt = y')$$

$$\Rightarrow y' = 50 - \frac{y}{100} = \frac{1}{100}(5,000 - y)$$

$$1 \text{ gal} = 3.78 \text{ l}$$

$$1 \text{ lb} = 0.454 \text{ kg}$$


► The initial condition :

$$y(0) = 100$$

1.3 Separable ODEs. Modeling

Step 1 Setting up a model.

$$\Rightarrow y' = 50 - \frac{y}{100} = \frac{1}{100}(5,000 - y) \quad y(0) = 100$$



Step 2 Solution of the model.


- General solution :

$$\frac{dy}{y-5,000} = -\frac{1}{100} dt \quad \Rightarrow \quad \ln|y-5,000| = -\frac{1}{100}t + c^* \quad \Rightarrow \quad y-5,000 = ce^{-\frac{t}{100}}$$

- Particular solution :

$$y(0) = 5,000 + ce^0 = 5,000 + c = 100 \quad \Rightarrow \quad c = -4,900$$

$$y = 5,000 - 4,900 e^{-\frac{t}{100}}$$



1.3 Separable ODEs. Modeling

☑ Newton's Law of Cooling


- The rate at which the temperature change of a body (물) is proportional to the difference between temperature of the body and the temperature of the surrounding medium.

$$\frac{dT}{dt} \propto (T - T_A) \Rightarrow \frac{dT}{dt} = k(T - T_A)$$

where, $k < 0$

T : Body temperature

T_A : Surrounding medium temperature





1.3 Separable ODEs. Modeling

✓ Newton's Law of Cooling

Great Idea !!

Relation between
 $\frac{dT(t)}{dt}$ and $T - T_A$.



T : body temperature
 T_A : outside temperature (constant)
 T_1, T_2 : Initial Body temperatures

$T_1 > T_A$, $T_2 < T_A$

1.3 Separable ODEs. Modeling

☑ Newton's Law of Cooling

$$\frac{dT}{dt} = k(T - T_A), \quad k < 0$$

$$\frac{dT}{T - T_A} = k \cdot dt$$

$$Y = T - T_A$$

$$\frac{dY}{dT} = 1 \Rightarrow dY = dT$$

$$\frac{dT}{T - T_A} = k \cdot dt$$

$$\frac{dY}{Y} = k \cdot dt$$

$$\int \frac{dY}{Y} = \int k \cdot dt$$

$$\ln|Y| + c_L = kt + c_R$$

$$\begin{aligned} \ln|Y| &= kt + c_R - c_L \\ &= kt + c \end{aligned}$$

1.3 Separable ODEs. Modeling

☑ Newton's Law of Cooling

①

$$\frac{dT}{dt} = k(T - T_A),$$
$$k < 0$$

$$\ln|Y| = kt + c$$

$$e^{\ln|Y|} = e^{kt+c}$$

$$|Y| = e^{kt+c}$$

②

$$|Y| = \tilde{c}e^{kt}$$

If $Y \geq 0$,

$$Y = \tilde{c}e^{kt}$$

If $Y < 0$,

$$-Y = \tilde{c}e^{kt}$$

$$Y = -\tilde{c}e^{kt}$$

③

$$Y = Ce^{kt}$$

, ($C = \text{sgn}(Y = T - T_A) \cdot \tilde{c}$)

$$T - T_A = Ce^{kt}$$

$$T = Ce^{kt} + T_A$$

$$\frac{dT}{dt} = k(T - T_A)$$

1.3 Separable ODEs. Modeling

- ☑ **Ex. 6** Suppose that in winter the daytime temperature in a certain office building is maintained at 70°F → **Initial condition**
- The heating is **shut off at 10 P.M.** and **turned on again at 6 A.M.**
 - On a certain day **the temperature inside the building = 65°F at 2 A.M.**
 - The outside temperature: **50°F at 10 P.M.** ~ **40°F by 6 A.M.**

What was **the temperature inside the building (T)** when the heat was turned on at 6 A.M.? _____

Step 1 Setting up a model

Temperature inside the building $T(t)$, Outside temperature T_A

$$\frac{dT}{dt} = k(T - T_A)$$

Step 2 General Solution

T_A varied between **50°F to 40°F**,

Golden Rule: If you cannot solve your problem, try to solve a simpler one.

$$T_A = 45^\circ\text{F} \quad \frac{dT}{(T - 45)} = k dt \quad T(t) = 45 + Ce^{kt}$$

Step 3 Particular solution Let 10 P.M to $t=0$. → $T(0)=70$

$$T(0) = 45 + Ce^0 = 70 \Rightarrow C = 25, \quad T_p(t) = 45 + 25e^{kt}$$

1.3 Separable ODEs. Modeling

- ☑ **Ex. 6** Suppose that in winter the daytime temperature in a certain office building is maintained at 70°F.
- The heating is shut off at 10 P.M. and turned on again at 6 A.M.
 - On a certain day the temperature inside the building = 65°F at 2 A.M.
 - The outside temperature: 50°F at 10 P.M. ~ 40°F by 6 A.M.

What was the temperature inside the building (T) when the heat was turned on at 6 A.M.?


Step 4 Determination of k $T(4)=65$

$$T_p(4) = 45 + 25e^{4k} = 65 \quad e^{4k} = 0.8 \quad k = \frac{1}{4} \ln 0.8 = -0.056$$

$$T_p(t) = 45 + 25e^{-0.056t}$$

Step 5 Answer and interpretation 6 A.M is $t=8$

$$T_p(8) = 45 + 25e^{-0.056 \cdot 8} = 61[^\circ \text{F}]$$



Particular solution (temperature) in Example 6

1.3 Separable ODEs. Modeling

❖ **Extended Method (확장방법)** : Reduction to Separable Form. Certain first order equations that are not separable can be made separable by a simple change of variables.

- A homogeneous ODE $y' = f\left(\frac{y}{x}\right)$ can be reduced to separable form by the substitution of $y=ux$

$$y' = f\left(\frac{y}{x}\right) \Rightarrow u'x + u = f(u) \Rightarrow \frac{du}{f(u)-u} = \frac{dx}{x} \quad \left(y = ux \Rightarrow u = \frac{y}{x} \text{ \& } y' = (ux)' = u'x + u \right)$$

Q?

- ☑ **Ex. 8** Solve $2xyy' = y^2 - x^2$

1.4 Exact ODEs, Integrating Factors

1.4 Exact ODEs, Integrating Factors

❖ Exact Differential Equation (완전미분 방정식):

The ODE $M(x, y)dx + N(x, y)dy = 0$ whose the differential form $M(x, y)dx + N(x, y)dy$ is **exact** (완전미분), that is, this form is the differential $du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$ of $u(x, y)$

$M(x, y)$ $N(x, y)$

- If ODE is an exact differential equation, then

$$M(x, y)dx + N(x, y)dy = 0 \quad \Rightarrow \quad du = 0 \quad \Rightarrow \quad u(x, y) = c$$

- ❖ Condition for exactness: $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ $\left(\because \frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(\frac{\partial u}{\partial x} \right) = \frac{\partial^2 u}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) = \frac{\partial N}{\partial x} \right)$

- ❖ Solve the exact differential equation.

$$M(x, y) = \frac{\partial u}{\partial x} \quad \Rightarrow \quad u(x, y) = \int M(x, y) dx + k(y) \quad \Rightarrow \quad \frac{\partial u}{\partial y} = N(x, y) \quad \Rightarrow \quad \frac{dk}{dy} \quad \& \quad k(y)$$

$$N(x, y) = \frac{\partial u}{\partial y} \quad \Rightarrow \quad u(x, y) = \int N(x, y) dy + l(x) \quad \Rightarrow \quad \frac{\partial u}{\partial x} = M(x, y) \quad \Rightarrow \quad \frac{dl}{dx} \quad \& \quad l(x)$$

1.4 Exact ODEs, Integrating Factors

✓ **Ex. 1 Solve** $\cos(x+y)dx + (3y^2 + 2y + \cos(x+y))dy = 0$

Step 1 Test for exactness.

$$\begin{array}{l} M(x, y) = \cos(x+y) \Rightarrow \frac{\partial M}{\partial y} = -\sin(x+y) \\ N(x, y) = 3y^2 + 2y + \cos(x+y) \Rightarrow \frac{\partial N}{\partial x} = -\sin(x+y) \end{array} \quad \left. \vphantom{\begin{array}{l} M(x, y) = \cos(x+y) \\ N(x, y) = 3y^2 + 2y + \cos(x+y) \end{array}} \right\} \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Step 2 Implicit general solution.

$$\begin{aligned} u(x, y) &= \int M(x, y)dx + k(y) = \int \cos(x+y)dx + k(y) = \sin(x+y) + k(y) \\ \Rightarrow \frac{\partial u}{\partial y} &= \cos(x+y) + \frac{dk}{dy} = N(x, y) \Rightarrow \frac{dk}{dy} = 3y^2 + 2y \Rightarrow k = y^3 + y^2 + c^* \end{aligned}$$

$$\therefore u(x, y) = \sin(x+y) + y^3 + y^2 = c$$

Step 3 Checking an implicit solution.

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$$

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy = \cos(x+y)dx + (\cos(x+y) + 3y^2 + 2y)dy = 0$$

1.4 Exact ODEs, Integrating Factors

✓ Example Q?

Solving an Exact DE

Solve $2xydx + (x^2 - 1)dy = 0$

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$$

$M(x, y)$ $N(x, y)$

1.4 Exact ODEs, Integrating Factors

❖ Reduction to Exact Form, Integrating Factors (적분 인자)

Some equations can be made exact by multiplication by some function, $F(x, y) \neq 0$, which is usually called the Integrating Factor.

☑ Ex. 3 Breakdown in the Case of Nonexactness

$$-ydx + xdy = 0$$

$$\because \frac{\partial}{\partial y}(-y) = -1, \quad \frac{\partial}{\partial x}(x) = 1 \quad \Rightarrow \quad \text{That equation is not exact.}$$

If we multiply it by $\frac{1}{x^2}$, we get an exact equation

$$-\frac{y}{x^2} dx + \frac{1}{x} dy = 0 \left(\because \frac{\partial}{\partial y} \left(-\frac{y}{x^2} \right) = -\frac{1}{x^2} = \frac{\partial}{\partial x} \left(\frac{1}{x} \right) \right)$$

General solution $y/x = c$

1.4 Exact ODEs, Integrating Factors

❖ How to Find Integrating Factors (F) ?

$$FPdx + FQdy = 0$$

The exactness condition : $\frac{\partial}{\partial y}(FP) = \frac{\partial}{\partial x}(FQ) \Rightarrow \frac{\partial F}{\partial y}P + F\frac{\partial P}{\partial y} = \frac{\partial F}{\partial x}Q + F\frac{\partial Q}{\partial x}$

Golden Rule : If you cannot solve your problem, try to solve a simpler one.

Hence we look for an integrating factor depending only on one variable.

Case 1)

$$F = F(x) \Rightarrow \frac{\partial F}{\partial x} = F', \quad \frac{\partial F}{\partial y} = 0 \quad \text{“}F=F(x,y) \text{ 가 일반적이지만, 단순히 } F(x) \text{로 가정”}$$

$$FP_y = F'Q + FQ_x$$

$$F'Q = F(P_y - Q_x)$$

$$\frac{F'}{F} = \frac{1}{Q}(P_y - Q_x)$$

$$\frac{1}{F} \frac{dF}{dx} = R(x) \quad \text{where } R(x) = \frac{1}{Q} \left(\frac{\partial P}{\partial y} - \frac{\partial Q}{\partial x} \right)$$

$$\therefore F(x) = \exp\left(\int R(x)dx\right)$$

Q?

Case 2)

1.4 Exact ODEs, Integrating Factors

☑ **Ex.** Find an integrating factor and solve the initial value problem

$$(e^{x+y} + ye^y)dx + (xe^y - 1)dy = 0, \quad y(0) = -1$$

Step 1 Nonexactness.

$$\begin{aligned} P(x, y) = e^{x+y} + ye^y &\Rightarrow \frac{\partial P}{\partial y} = e^{x+y} + e^y + ye^y \\ Q(x, y) = xe^y - 1 &\Rightarrow \frac{\partial Q}{\partial x} = e^y \end{aligned} \quad \left. \vphantom{\begin{aligned} P(x, y) = e^{x+y} + ye^y \\ Q(x, y) = xe^y - 1 \end{aligned}} \right\} \frac{\partial P}{\partial y} \neq \frac{\partial Q}{\partial x}$$

$$\begin{aligned} R(x) &= \frac{1}{Q} \left(\frac{\partial P}{\partial y} - \frac{\partial Q}{\partial x} \right) \\ F(x) &= \exp\left(\int R(x) dx\right) \end{aligned}$$

$$\begin{aligned} R^*(y) &= \frac{1}{P} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \\ F^*(y) &= \exp\left(\int R^*(y) dy\right) \end{aligned}$$

Step 2 Integrating factor. General solution.

$$R^*(y) = \frac{1}{P} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) = \frac{1}{e^{x+y} + ye^y} (e^y - e^{x+y} - e^y - ye^y) = -1 \Rightarrow F^*(y) = e^{-y}$$

$$\therefore (e^x + y)dx + (x - e^{-y})dy = 0 \quad \text{is the exact equation.}$$

Q? Why not R?

1.4 Exact ODEs, Integrating Factors

✓ Ex. Find an integrating factor and solve the initial value problem

$$(e^{x+y} + ye^y)dx + (xe^y - 1)dy = 0, \quad y(0) = -1$$

Step 2 Integrating factor. General solution. $F^*(y) = e^{-y}$

$$M(x, y) = \frac{\partial u}{\partial x}$$

$$u(x, y) = \int M(x, y) dx + k(y)$$

$$\frac{\partial u}{\partial y} = N(x, y)$$

$$\underbrace{(e^x + y)}_{M(x, y)} dx + \underbrace{(x - e^{-y})}_{N(x, y)} dy = 0$$

$$u = \int (e^x + y) dx = e^x + xy + k(y)$$

$$\Rightarrow \frac{\partial u}{\partial y} = x + k'(y) = x - e^{-y} \quad \Rightarrow \quad k'(y) = -e^{-y}, \quad k(y) = e^{-y}$$

The general solution is $u(x, y) = e^x + xy + e^{-y} = c$

Step 3 Particular solution

$$y(0) = -1 \quad \Rightarrow \quad u(0, -1) = e^0 + 0 + e = 3.72$$

$$\therefore u(x, y) = e^x + xy + e^{-y} = 3.72$$

1.4 Exact ODEs, Integrating Factors

☑ Example

Nonexact ODE


Solve

Q? Integration
Factor

$$xydx + (2x^2 + 3y^2 - 20)dy = 0$$

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics



❖ **Linear ODEs:** ODEs which is linear in both the unknown function (y) and its derivative (y').

Ex. $y' + p(x)y = r(x)$: Linear differential equation

$y' + p(x)y = r(x)y^2$: Nonlinear differential equation

■ Standard Form : $y' + p(x)y = r(x)$ ($r(x)$: Input, $y(x)$: Output)

❖ **Homogeneous, Nonhomogeneous Linear ODE**

$y' + p(x)y = 0$: Homogeneous Linear ODE

$y' + p(x)y = r(x) \neq 0$: Nonhomogeneous Linear ODE

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

- ❖ **Homogeneous Linear ODE** (Apply the method of separating variables)

$$y' + p(x)y = 0 \quad \Rightarrow \quad y = ce^{-\int p(x)dx} \quad (1^*)$$

- ❖ **Nonhomogeneous Linear ODE** (Find integrating factor and solve)

$$y' + p(x)y = r(x) \quad \Rightarrow \quad (py - r)dx + dy = 0 \quad \text{is not exact}$$

$$\left(\because \frac{\partial}{\partial y}(py - r) = p \neq 0 = \frac{\partial}{\partial x}(1) \right)$$

- Find integrating factor. We multiply $F(x)$.

$$Fy' + pFy = rF \quad \Rightarrow \quad \text{If } pFy = F'y \quad \Rightarrow \quad Fy' + F'y = rF$$

$$F(x)y = \left(\int r(x)F(x)dx + c \right)$$

$$\Rightarrow (Fy)' = rF \quad \Rightarrow \quad y = \frac{1}{F(x)} \left(\int r(x)F(x)dx + c \right)$$

- 즉, $pF = F'$ 이 되는 F 를 찾아서 양변에 곱하면 \rightarrow **Exact ODE**

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

❖ **Nonhomogeneous Linear ODE** (Find integrating factor and solve)

$$y' + p(x)y = r(x) \Rightarrow (py - r)dx + dy = 0 \quad \text{is not exact} \quad \left(\because \frac{\partial}{\partial y}(py - r) = p \neq 0 = \frac{\partial}{\partial x}(1) \right)$$

From exactness condition

$$Fy' + pFy = rF \quad \rightarrow \quad Fdy + (pFy - rF) dx = 0$$

$$\rightarrow (pFy - rF)dx + Fdy = 0$$

$$\rightarrow \frac{\partial}{\partial y}(pFy - rF) = \frac{\partial F}{\partial x}$$

$$\rightarrow \mathbf{pF = F'}$$

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

- ❖ **Nonhomogeneous Linear ODE** (Find integrating factor and solve)

$$y' + p(x)y = r(x) \Rightarrow (py - r)dx + dy = 0$$

- Find integrating factor (F) from $pF = F'$

- By separating variables,

$$p = \frac{F'}{F} \Rightarrow p = \frac{dF}{dx} \frac{1}{F} \Rightarrow p dx = \frac{dF}{F}$$

- By integration, writing $h = \int p dx$,

$$\ln |F| = h = \int p dx, \Rightarrow F = e^h$$

- With $F = e^h$ and $h' = p$, Eq. $Fy' + pFy = rF$ becomes

$$e^h y' + h' e^h y = e^h y' + (e^h)' y = \boxed{(e^h y)'} = r e^h \Rightarrow e^h y = \int e^h r dx + c$$

- By integration,

$$y(x) = e^{-h} \left(\int e^h r dx + c \right) = e^{-h} \int e^h r dx + c e^{-h}, \quad h = \int p(x) dx$$

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

☑ Ex. 1 Solve the linear ODE $y' - y = e^{2x}$

$$\frac{dy}{dx} - (y + e^{2x}) = 0 \Rightarrow (y + e^{2x})dx - dy = 0$$

$$(y + e^{2x})Fdx - Fdy = 0, \quad F = F(x)$$

$$\text{Exactness : } \frac{\partial}{\partial y}(yF + e^{2x}F) = -\frac{\partial F}{\partial x} \Rightarrow F = -F'$$

$(pF = F' \Rightarrow p = -1)$

$$\frac{dF}{F} = -dx \Rightarrow \ln F = -x \Rightarrow F = e^{-x}$$

$$(y + e^{2x})e^{-x}dx - e^{-x}dy = 0 \quad (Fy' + pFy = rF)$$

$$\text{where, } \frac{\partial}{\partial y}(ye^{-x} + e^x) = e^{-x}, \quad \frac{\partial}{\partial x}(-e^{-x}) = e^{-x}$$

$$\rightarrow e^{-x}y' - e^{-x}y = e^{2x}e^{-x} \Rightarrow (e^{-x}y)' = e^x$$

$$(e^{-x}y)dy = e^x dx \Rightarrow e^{-x}y = e^x + c$$

$$y = e^{2x} + ce^x$$

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

☑ Ex. 1 Solve the linear ODE $y' - y = e^{2x}$

$$y' + p(x)y = r(x)$$

$$y(x) = e^{-h} \left(\int e^h r dx + c \right), \quad h = \int p(x) dx$$
$$= e^{-h} \int e^h r dx + ce^{-h}$$

$$p = -1, \quad r = e^{2x}, \quad h = \int p dx = -x \quad \Rightarrow \quad \therefore y = e^{-h} \left[\int e^h r dx + c \right] = e^x \left[\int e^{-x} e^{2x} dx + c \right] = e^x \left[e^x + c \right] = e^{2x} + ce^x$$

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

❖ **Bernoulli Equation:** $y' + p(x)y = g(x)y^a$ ($a \neq 0$ & 1): Nonlinear ODE

We set $u(x) = [y(x)]^{1-a}$

$$y' = g(x)y^a - p(x)y$$

$$\Rightarrow u' = (1-a)y^{-a} \boxed{y'} = (1-a)y^{-a}(gy^a - py) = (1-a)(g - py^{1-a}) = (1-a)(g - pu)$$

$$\Rightarrow u' + (1-a)pu = (1-a)g$$

: Now transformed to Linear ODE

$$\boxed{y' + p(x)y = r(x)}$$

$$\boxed{y(x) = e^{-h} \left(\int e^h r dx + c \right), \quad h = \int p(x) dx}$$

1.5 Linear ODEs. Bernoulli Equation. Population Dynamics

❖ **Bernoulli Equation:** $y' + p(x)y = g(x)y^a$ ($a \neq 0$ & 1)

We set $u(x) = [y(x)]^{1-a}$

☑ Ex. 4 Logistic Equation

Solve the following Bernoulli equation, known as the logistic equation (or Verhulst equation) $y' = Ay - By^2$

$$y' = Ay - By^2 \Rightarrow y' - Ay = -By^2 \quad \& \quad a = 2 \quad (u = y^{-1})$$

$$\Rightarrow u' = -y^{-2}y' = -y^{-2}(Ay - By^2) = -Ay^{-1} + B = -Au + B \Rightarrow u' + Au = B$$

$$p = A, r = B \Rightarrow h = \int p dx = Ax \quad \& \quad u = e^{-h} \left[\int e^h r dx + c \right] = e^{-Ax} \left[\frac{B}{A} e^{Ax} + c \right] = ce^{-Ax} + \frac{B}{A}$$

$$y(x) = e^{-h} \left(\int e^h r dx + c \right), \quad h = \int p(x) dx$$

The general solution of the equation is $y = \frac{1}{u} = \frac{1}{\left(\frac{B}{A} + ce^{-Ax}\right)}$

1.6 Orthogonal Trajectories (직교 절선) - Skip

❖ Orthogonal Trajectory

: A family of curves in the plane that intersect a given family of curves at given angles.

❖ Find the orthogonal trajectories by using ODEs.

Step 1 Find an ODE $y' = f(x, y)$ for which the give family is a general solution.

Step 2 Write down the ODE $y' = -\frac{1}{f(x, y)}$ of the orthogonal trajectories.

Step 3 Solve it.

☑ Ex. A one-parameter family of quadratic parabolas is given by $y = cx^2$ 

Step 1 $\frac{y}{x^2} = c \Rightarrow \frac{y'x^2 - 2xy}{x^4} = 0 \Rightarrow y' = \frac{2y}{x}$

Step 2 $y' = -\frac{x}{2y}$

Step 3 $2yy' + x = 0 \Rightarrow y^2 + \frac{1}{2}x^2 = c^*$

1.7 Existence and Uniqueness of Solutions for Initial Value Problems

1.7 Existence and Uniqueness of Solutions for Initial Value Problems

- ❖ An initial value problem may have no solution, precisely one solution, or more than one solution.
- Ex. $|y'| + |y| = 0, \quad y(0) = 1 \quad \Rightarrow \quad$ No solution
- $y' = 2x, \quad y(0) = 1 \quad \Rightarrow \quad$ Precisely one solution $\Rightarrow y = x^2 + 1$
- $xy' = y - 1, \quad y(0) = 1 \quad \Rightarrow \quad$ Infinitely many solutions $\Rightarrow y = 1 + cx$

❖ Problem of Existence (존재성)

Under what conditions does an initial value problem have at least one solution (hence one or several solutions)?

❖ Problem of Uniqueness (유일성)

Under what conditions does that problem have at most one solution (hence excluding the case that has more than one solution)?

1.7 Existence and Uniqueness of Solutions for Initial Value Problems

❖ Theorem 1 Existence Theorem (존재 정리)

Let the right side $f(x,y)$ of the ODE in the initial value problem.

$$(1) \quad y' = f(x, y), \quad y(x_0) = y_0$$


be continuous at all points (x, y) in some rectangle

$$R : |x - x_0| < a, \quad |y - y_0| < b$$

and bounded in R ; that is, there is a number K such that

$$(2) \quad |f(x, y)| \leq K \quad \text{for all } (x, y) \text{ in } R.$$

Then the initial value problem (1) has at least one solution $y(x)$. This solution exists at least for all x in the subinterval $|x - x_0| < \alpha$ of the interval $|x - x_0| < a$; here, α is the smaller of the two numbers a and b/K .



EX) $f(x, y) = x^2 + y^2$ is bounded (with $K = 2$) in the square of $|x| < 1, |y| < 1$.

$f(x, y) = \tan(x+y)$ is not bounded for $|x+y| < \pi/2$

1.7 Existence and Uniqueness of Solutions for Initial Value Problems

❖ Theorem 2 Uniqueness Theorem (유일성 정리)


Let f and its partial derivative $f_y = \partial f / \partial y$ be continuous for all (x, y) in the rectangle R and bounded, say,

(3) (a) $|f(x, y)| \leq K$ (b) $|f_y(x, y)| \leq M$ for all (x, y) in R .


Then the initial value problem (1) has at most one solution $y(x)$. Thus, by the Existence Theorem, the problem has precisely one solution. This solution exists at least for all x in that subinterval $|x - x_0| < \alpha$.

$y' = f(x, y)$, the condition (2) implies that $|y'| \leq K$

Case of $b \geq aK$
 $\Rightarrow \alpha = a$
 Solution exists for
 $x_0 - a \leq x \leq x_0 + a$



Case of $b < aK$
 $\Rightarrow \alpha = b / K < a$
 Solution exists for
 $x_0 - b / K \leq x \leq x_0 + b / K$



1.7 Existence and Uniqueness of Solutions for Initial Value Problems

☑ **Ex. 1** Consider initial value problem

$$y' = 1 + y^2, \quad y(0) = 0$$

R ; $|x| < 5$, $|y| < 3$, then, $a = 5, b = 3$ and

$$|f(x, y)| = |1 + y^2| \leq K = 10$$

$$\left| \frac{\partial f}{\partial y} \right| = 2|y| \leq M = 6 \Rightarrow \alpha = \frac{b}{K} = 0.3 < a$$

The solution of the problem $y = \tan x$. It is discontinuous at $\pm\pi/2$ and no continuous solution valid in the entire interval from which we started $|x| < 5$.

$$\begin{aligned} \frac{y'}{1+y^2} = 1 &\Rightarrow \frac{dy/dx}{1+y^2} = 1 &\Rightarrow \frac{dy}{1+y^2} = dx \\ &\Rightarrow \int \frac{1}{1+y^2} dy = \int dx + c &\Rightarrow \arctan y = x + c &\Rightarrow y = \tan(x + c) \\ &\Rightarrow y = \tan(x + c) \quad (\because y(0) = 0) \end{aligned}$$

[Reference] Natural Logarithm (ln(x)) function

Rule name	Rule	Example
Product rule	$\ln(x \cdot y) = \ln(x) + \ln(y)$	$\ln(3 \cdot 7) = \ln(3) + \ln(7)$
Quotient rule	$\ln(x / y) = \ln(x) - \ln(y)$	$\ln(3 / 7) = \ln(3) - \ln(7)$
Power rule	$\ln(x^y) = y \cdot \ln(x)$	$\ln(2^8) = 8 \cdot \ln(2)$
In derivative	$f(x) = \ln(x) \Rightarrow f'(x) = 1/x$	
In integral	$\int \ln(x) dx = x \cdot (\ln(x) - 1) + C$	
In of negative number	$\ln(x)$ is undefined when $x \leq 0$	
In of zero	$\ln(0)$ is undefined	
	$\lim_{x \rightarrow 0^+} \ln(x) = -\infty$	
In of one	$\ln(1) = 0$	
In of infinity	$\lim \ln(x) = \infty$, when $x \rightarrow \infty$	