







Overview on Scattering




Scattering pattern of nanocrystal



$$I(q) = I_0 N (\rho - \rho_0)^2 F^2(q)$$

$$F(q) = \frac{4}{3} \pi R^3 \left[3 \frac{\sin(qR) - qR \cos(qR)}{(qR)^3} \right]$$

elastic scattering: $k_i = k_f$



for constructive interference: $2\vec{k} \cdot \vec{s} = |\vec{k}|^2$

$$n\lambda = 2d \sin \theta$$

Properties of x-rays and neutrons

x-rays:

electromagnetic
radiation

$$c = \lambda \nu$$

$$E = h\nu$$

$$p = h / \lambda$$

$$1/\nu = \tau = 10^{-19} \text{ s}$$

neutron:

an uncharged
elementary particle

ν depends on λ


$$E = m\nu^2/2$$

$$p = m\nu$$

$$1/\nu = \tau = 10^{-13} \text{ s}$$

magnetic moment

v distribution from reactor:-dep. on moderator Maxwell-Boltzmann distribution



$-kT$ (at RT) ~ 20 meV

investigation on dynamics is also available.

$$J = |A|^2 = AA^*$$


$$\frac{J}{J_0} = \frac{d\sigma}{d\Omega}$$

- differential scattering cross-section:
the probability that a photon or a neutron impinging on the sample into a unit solid angle in the given direction


$$\frac{J}{J_0} = \frac{d\sigma}{d\Omega}$$

$$= \frac{\text{number of particles scattered into a unit solid angle per second}}{\text{flux of incident beam}}$$

$$\sigma_{\text{tot}} = \frac{\text{total number of particles scattered in all direction per second}}{\text{flux of incident beam}}$$




elastic scattering: $k_i = k_f$



for constructive interference: $2\vec{k} \cdot \vec{s} = |\vec{k}|^2$

$$n\lambda = 2d \sin \theta$$


elastic scattering: $k_i = k_f$



for constructive interference: $2\vec{k} \cdot \vec{s} = |\vec{k}|^2$


$$n\lambda = 2d \sin \theta$$

coherent scattering:



$$\Delta\phi = \frac{2\pi\delta}{\lambda} = \frac{2\pi}{\lambda} (\vec{S}_0 \cdot \vec{r} - \vec{S} \cdot \vec{r})$$
$$= -2\pi\vec{s} \cdot \vec{r}$$

$$\vec{s} = \frac{\vec{S} - \vec{S}_0}{\lambda}$$



at the detector:

$$A = A_1 + A_2 = A_0^2 b^2 e^{i2\pi(vt-x/\lambda)} (1 + e^{-i2\pi\vec{s}\cdot\vec{r}})$$

at the detector:

$$A = A_1 + A_2 = A_0^2 b^2 e^{i2\pi(vt-x/\lambda)} (1 + e^{-i2\pi\vec{s}\cdot\vec{r}})$$

$$J = AA^* = A_0^2 b^2 (1 + e^{i2\pi\vec{s}\cdot\vec{r}})(1 + e^{-i2\pi\vec{s}\cdot\vec{r}})$$

$$A = A_0 b (1 + e^{-i2\pi\vec{s}\cdot\vec{r}})$$

when there are N identical scatterers,

$$A = A_0 b \sum_{j=1}^N e^{-i2\pi\vec{s}\cdot\vec{r}_j} = A_0 b \sum_{j=1}^N e^{-i\vec{q}\cdot\vec{r}_j}$$

$$A = A_0 b \int_V n(\vec{r}) e^{-i\vec{q}\cdot\vec{r}} d\vec{r}$$

Real and inverse lattice

$$\vec{r} = u_1 \vec{a}_1 + u_2 \vec{a}_2 + u_3 \vec{a}_3$$

$$\vec{q} = v_1 \vec{b}_1 + v_2 \vec{b}_2 + v_3 \vec{b}_3$$

$$b_1 = 2\pi \frac{a_2 \times a_3}{a_2 \cdot a_3 \times a_1}$$

$$b_2 = 2\pi \frac{a_3 \times a_1}{a_3 \cdot a_1 \times a_2}$$


$$b_3 = 2\pi \frac{a_1 \times a_2}{a_1 \cdot a_2 \times a_3}$$


$$b_i \cdot a_j = 2\pi \delta_{ij}$$

when $i \neq j$, 0




when $i = j$, 2π

$$2\pi \vec{s} = \vec{q}$$





unit cell from a certain angle

-  real lattice
-  diffracted spot
-  beam center

Structure factor or form factor


$$A(q) \sim F(q) \quad \rho(\vec{r}) = \sum_j \rho_j(\vec{r} - \vec{r}_j)$$

$$F(q) = \int \rho(\vec{r}) \exp(-i\vec{r} \cdot \vec{q}) d\vec{r}$$

$$= \sum_j \int \rho(\vec{r} - \vec{r}_j) \exp(-i\vec{r} \cdot \vec{q}) d\vec{r}$$

$$= \sum_j \left[\int_V \rho(\vec{r} - \vec{r}_j) \exp[-i\vec{q} \cdot (\vec{r} - \vec{r}_j)] d\vec{r} \right] \exp(-i\vec{q} \cdot \vec{r}_j)$$

$$= \sum_j f_j \exp(-i\vec{q} \cdot \vec{r}_j)$$



Atomic form factor:
intensity determination

Position determination

Structure factor or form factor

$$A(q) \sim F(q) \quad \rho(\vec{r}) = \sum_j \rho_j(\vec{r} - \vec{r}_j)$$

$$F(q) = \int \rho(\vec{r}) \exp(-i\vec{r} \cdot \vec{q}) d\vec{r}$$

assuming spherical symmetry,

$$= \int 4\pi r^2 \rho(r) \frac{\sin(\vec{r} \cdot \vec{q})}{\vec{r} \cdot \vec{q}} dr$$

see Fig 1.6

- Scattering length of a single nucleus

- interaction w/ nucleus:

 - highly penetrating

- scattering occurs due to

1. structure

2. randomness of spin state

 - and distribution of isotope

- Coherent and incoherent scattering length

$$i + 1/2 \quad 2(i + 1/2) + 1 = 2i + 2 \quad f^+ = \frac{2i + 2}{4i + 2} = \frac{i + 1}{2i + 1}$$

$$i - 1/2 \quad 2(i - 1/2) + 1 = 2i \quad f^- = \frac{2i}{4i + 2} = \frac{i}{2i + 1}$$


Ewald sphere and reciprocal scattering

$$A(\vec{q}) = \int \rho(\vec{r}) e^{-i\vec{r}\cdot\vec{q}} d\vec{r}$$

SLDD

Scattering amplitude

$$\rho(\vec{r}) = \rho_u(\vec{r}) * z(\vec{r})$$



$$A(\vec{q}) = F(\vec{q})Z(\vec{q})$$

Form factor

lattice factor

autocorrelation

$$\begin{aligned} \Gamma_\rho(\vec{r}) &= V \langle \rho(\vec{u}) \rho(\vec{u}') \rangle \\ &= \int \rho(\vec{r}) \rho(\vec{u} + \vec{r}) d\vec{u} \end{aligned}$$




squaring

$$\begin{aligned} I(\vec{q}) &= A(\vec{q}) \cdot A^*(\vec{q}) \end{aligned}$$

$$I(\vec{q})$$

$$\Gamma_\rho(\vec{r})$$

Autocorrelation ftn



$$I(\vec{q}) = \int \Gamma_\rho(\vec{r}) e^{-i\vec{r}\cdot\vec{q}} d\vec{r}$$

$$A(\vec{q}) = \int \rho(\vec{r}) e^{-i\vec{r} \cdot \vec{q}} d\vec{r}$$

SLDD

$$\rho(\vec{r}) = \rho_u(\vec{r}) * z(\vec{r})$$

FT

Scattering amplitude

$$A(\vec{q}) = F(\vec{q})Z(\vec{q})$$

IFT

Form factor

lattice factor



squaring

$$I(\vec{q})$$

$$= A(\vec{q}) \cdot A^*(\vec{q})$$



$$I(\vec{q})$$

SLDD


$$\rho(\vec{r}) = \rho_u(\vec{r}) * z(\vec{r}) \xrightarrow{\text{FT}} A(\vec{q}) = F(\vec{q})Z(\vec{q}) \xrightarrow{\text{IFT}}$$

Scattering amplitude

Form factor lattice factor



$$\rho_u(\vec{r}) \times z(\vec{r})$$



$$\rho(\vec{r}) = \rho_u(\vec{r}) * z(\vec{r})$$

$$\rho(x) * z(x)$$


$$= \sum_{n=-\infty}^{\infty} \rho(x) * \delta(x - na)$$

$$= \sum_{n=-\infty}^{\infty} \int_0^{x-na} \rho(u) \delta((x - na) - u) du$$

$$\left[\text{since } \int_0^{x-na} \rho(u) \delta((x - na) - u) du = f(x - na), \right]$$

$$= \sum_{n=-\infty}^{\infty} \rho(x - na)$$

$$\begin{aligned}\rho(\vec{r}) &= \rho_u(\vec{r}) * z(\vec{r}) \\ &= \sum_{n=-\infty}^{\infty} \rho(x - na)\end{aligned}$$



SLDD

$$\rho(\vec{r}) = \rho_u(\vec{r}) * z(\vec{r})$$

autocorrelation

$$\Gamma_\rho(\vec{r})$$

$$= V \langle \rho(\vec{u}) \rho(\vec{u}') \rangle$$


$$= \int \rho(\vec{r}) \rho(\vec{u} + \vec{r}) d\vec{u}$$




$$\Gamma_\rho(\vec{r})$$

Autocorrelation ftn

FT



IFT



$$I(\vec{q})$$

$$I(\vec{q}) = \int \Gamma_\rho(\vec{r}) e^{-i\vec{r} \cdot \vec{q}} d\vec{r}$$

$$I(\vec{q})$$

$$= \langle |A(\vec{q})|^2 \rangle = A(\vec{q}) \cdot A^*(\vec{q})$$

$$= \langle \left| \int \rho(\vec{r}) e^{i\vec{r} \cdot \vec{q}} d\vec{r} \right|^2 \rangle = A^*(\vec{q}) \cdot A(\vec{q})$$

$$= \left[\int \rho(\vec{u}) e^{i\vec{u} \cdot \vec{q}} d\vec{u} \right] \left[\int \rho(\vec{u}') e^{-i\vec{u}' \cdot \vec{q}} d\vec{u}' \right]$$

$$= \left[\int \rho(\vec{u}) e^{i\vec{u} \cdot \vec{q}} d\vec{u} \right] \left[\int \rho(\vec{r} + \vec{u}) e^{-i(\vec{r} + \vec{u}) \cdot \vec{q}} d\vec{r} \right]$$

$$= \int \left[\int \rho(\vec{u}) \rho(\vec{r} + \vec{u}) d\vec{u} \right] e^{-i\vec{r} \cdot \vec{q}} d\vec{r}$$


$$= \int \Gamma_\rho(\vec{r}) e^{-i\vec{r} \cdot \vec{q}} d\vec{r}$$

$$\vec{u}' = \vec{r} + \vec{u}$$

$$d\vec{u}' = d\vec{r}$$

Physical meaning of auto correlation ftn

$$\begin{aligned}\Gamma_{\rho}(\vec{r}) &= V \langle \rho(\vec{u}) \rho(\vec{u}') \rangle \\ &= \int \rho(\vec{r}) \rho(\vec{u} + \vec{r}) d\vec{u}\end{aligned}$$



see also p.96

$$\eta(\vec{r}) = \rho(\vec{r}) - \langle \rho \rangle$$

$$\Gamma_\rho(\vec{r})$$

$$= V \langle \rho(\vec{u}) \rho(\vec{u}') \rangle$$

$$= \int \rho(\vec{r}) \rho(\vec{u} + \vec{r}) d\vec{u}$$

$$= \int [\eta(\vec{u}) + \langle \rho \rangle] [\eta(\vec{u} + \vec{r}) + \langle \rho \rangle] d\vec{u}$$

$$= \int \eta(\vec{u}) \eta(\vec{u} + \vec{r}) d\vec{u} + \langle \rho \rangle^2 \int d\vec{u} + \langle \rho \rangle \int \eta(\vec{u}) d\vec{u} + \langle \rho \rangle \int \eta(\vec{u} + \vec{r}) d\vec{u}$$

$$\sim \int \eta(\vec{u}) \eta(\vec{u} + \vec{r}) d\vec{u} + \langle \rho \rangle^2 V$$

macroscopic dimension

=0

$$\sim \Gamma_\eta(\vec{r})$$


null scattering or scattering at $q=0$
 Experimentally unobservable

$$A(\vec{q}) = \int \rho(\vec{r}) e^{-i\vec{r}\cdot\vec{q}} d\vec{r}$$

SLDD

Scattering amplitude

$$\rho(\vec{r}) = \rho_u(\vec{r}) * z(\vec{r})$$



$$A(\vec{q}) = F(\vec{q})Z(\vec{q})$$

Form factor

lattice factor

autocorrelation


$$\Gamma_\rho(\vec{r}) = V \langle \rho(\vec{u}) \rho(\vec{u} + \vec{r}) \rangle = \int \rho(\vec{r}) \rho(\vec{u} + \vec{r}) d\vec{u}$$



squaring

$$I(\vec{q}) = A(\vec{q}) \cdot A^*(\vec{q})$$

$$\Gamma_\rho(\vec{r})$$



$$I(\vec{q})$$

Autocorrelation ftn


$$I(\vec{q}) = \int \Gamma_\rho(\vec{r}) e^{-i\vec{r}\cdot\vec{q}} d\vec{r}$$

Structure factor for a uniform sphere

$$2\pi\vec{s} = \vec{q}$$

$$\rho \sim 0$$

$$P(q) \sim [F(q)]^2$$



$$F(q) = \int \rho(\vec{r}) \exp(-i\vec{q} \cdot \vec{r}) d\vec{r}$$

= HOMEWORK!!! (by sep/22/2005)

Structure factors for several structures

Thin rod


At a certain orientations,

$$F(q) = \frac{4}{qL \cos \Theta} \sin\left(\frac{qL \cos \Theta}{2}\right)$$

Random orientations,

$$P(q) = \frac{2}{qL} \left[\int_0^{qL} \frac{\sin u}{u} du - \frac{1 - \cos qL}{qL} \right]$$

Homework!!!




Structure factors for several structures

Homework!!!


Circular disk

$$P(q) = \frac{2}{q^2 R^2} \left[1 - \frac{J_1(2qR)}{qR} \right]$$



Size of chain molecules

-synthetic polymer, DNA, protein...



Number of repeat unit

Charateristic ratio



$$R_g^2 = C_\infty n l^2 / 6$$

$$R_0^2 = 6R_g^2 = Nb^2$$

Number of Kuhn segment

Kuhn segment length

How about this?



$$P(q) \sim [F(q)]^2$$

constant form factor?

Random coil
Or Gaussian coil


$$P(q) \sim 2 \frac{\exp(-q^2 R_g^2) - 1 + q^2 R_g^2}{q^4 R_g^4}$$

Not only lattice scattering
but also shape of the single
object is important


$$I(q) = I_0 N (\rho - \rho_0)^2 F^2(q)$$


$$F(q) = \frac{4}{3} \pi R^3 \left[3 \frac{\sin(qR) - qR \cos(qR)}{(qR)^3} \right]$$

Log(Scattered Intensity) (arbitrary units)






Log(Scattered Intensity) (arbitrary units)





unit cell from a certain angle

-  real lattice
-  diffracted spot
-  beam center