

Chapter 1

Science, Technology, and Society

Contents

1

Technology and Society

2

Science and Technology

3

The Biology Century


Technology

- Try to imagine a world without technology.
 - Computer
 - TV
 - Car
 - Heating (Energy)
 - Cooling
 - Clothes
 - Food
 - House
 - Lamp, paper, pen, chair,

Technology and Society


- Technology
 - Changes the environment and society
 - Webster's definition:
“the totality of means employed to provide objects necessary for human sustenance and comfort”

Technology and Society



Technology and Society

- Society
 - Creates filters for technology



Science and Technology

■ Science

- Search for knowledge
- Way of understanding ourselves and the physical world
- Process of asking questions and finding answers, then creating broad generalizations
- Looks for order or patterns in the physical world
- Evaluated by how well the facts support the conclusion or theory
- Limited by the ability to collect relevant facts
- Discoveries give rise to technological advances

■ Technology

- Practical application of knowledge
- Way of adapting ourselves to the physical world
- Process of finding solutions to human problems to make lives easier and better
- Looks for ways to control the physical world
- Evaluated by how well it works
- Limited by financial costs and safety concerns
- Advances give rise to scientific discoveries

The Relationship Between Science and Technology

Technology

Experience-based

Provide tool
for scientific observation
and experiments

Science-based
technology


Science

Understanding of Nature


Derive technology
development

Derive progress
of science


Acceleration of
technological change



The Nature of Science



The Scientific Process



The Biology Century

- The past two centuries
 - Technology driven by physics and chemistry
 - Industrial Revolution, Information Age, Green Revolution
- The Biology Century will be fueled by biotechnology.

Biotechnology

- Definition
 - The use of living organisms or life processes to solve problems or make useful products
- Ancient biotechnology
 - Trial and error-based
- Modern biotechnology
 - The use of cells and biological molecules or cellular and biomolecular processes to solve problems and make useful products

Types and Applications of Biotechnology

Biotechnology

- Bioprocessing technol.
- Cell culture technol.
- Recombinant DNA technol.
(genetic engineering)
- Monoclonal antibody tech.
- Biosensor technol.
- Microarray technol.
- Protein engineering tech.



Industry


- Human health care
- Agricultural production
- Food and beverages
- Enzyme industry
- Chemical manufacturing
- Energy
- Waste treatment

Characteristics of Cells and Biomolecules

- Specificity, precision, and predictability
- Unity and flexibility
- Reproduction and renewable resources



The History of Crop Genetic Modification

- Stage1
 - Genetic modification through seed selection



The History of Crop Genetic Modification

- Stage 2
 - Genetic modification through plant breeding and selection
 - Invention of the microscope
 - Hand pollination




The History of Crop Genetic Modification

- Stage 3
 - Science-based plant breeding
 - Based on Mendel's work
- Stage 4
 - Plant genetic engineering


Model of Inheritance

- Fluid-blending model
- Discrete-particle model
 - Mendel's theory of inheritance (1865, proved in 1900)
 - Discrete-particle (now known as gene) model




The Nature of Genetic Material

- Frederick Griffith (1928)
 - 'Transforming factor' transferred from dead smooth virus to rough virus
- O.T. Avery (1943)
 - The 'transforming factor' was DNA



The Genetic Material Protein or DNA?

- The DNA-vs.-protein debate was resolved.
 - Alfred Hershey and Martha Chase (1952)
 - Identification of DNA as genetic material



DNA Double Helix

by J. Watson and F. Crick

NO. 4398 April 25, 1953 NATURE 737

MOLECULAR STRUCTURE OF NUCLEIC ACIDS

A Structure for Deoxyribose Nucleic Acid

WE wish to suggest a structure for the salt of deoxyribose nucleic acid (D.N.A.). This structure has novel features which are of considerable biological interest.

A structure for nucleic acid has already been proposed by Pauling and Corey¹. They kindly made their manuscript available to us in advance of publication. Their model consists of three intertwined chains, with the phosphates near the fibre axis, and the bases on the outside. In our opinion, this structure is unsatisfactory for two reasons: (1) We believe that the material which gives the X-ray diagram is the salt, not the free acid. Without the acidic hydrogen atoms it is not clear what forces would hold the structure together, especially as the negatively charged phosphates near the axis will repel each other. (2) Some of the van der Waals distances appear to be too small.

Another three-chain structure has also been suggested by Fraser (in the press). In his model the phosphates are on the outside and the bases on the inside, linked together by hydrogen bonds. This structure as described is rather ill-defined, and for this reason we shall not comment on it.

We wish to put forward a radically different structure for the salt of deoxyribose nucleic acid. This structure has two helical chains each coiled round the same axis (see diagram). We have made the usual chemical assumptions, namely, that each chain consists of phosphate diester groups joining β -D-deoxyribofuranose residues with 3',5' linkages. The two chains (but not their bases) are related by a dyad perpendicular to the fibre axis. Both chains follow right-handed helices, but owing to the dyad the sequences of the atoms in the two chains run in opposite directions. Each chain loosely resembles Furburg's² model No. 1; that is, the bases are on the inside of the helix and the phosphates on the outside. The configuration of the sugar and the atoms near it is close to Furburg's 'standard configuration', the sugar being roughly perpendicular to the attached base. There is a residue on each chain every 3.4 Å, in the z-direction. We have assumed an angle of 36° between adjacent residues in the same chain, so that the structure repeats after 10 residues on each chain, that is, after 34 Å. The distance of a phosphate atom from the fibre axis is 10 Å. As the phosphates are on the outside, cations have easy access to them.

The structure is an open one, and its water content is rather high. At lower water contents we would expect the bases to tilt so that the structure would become more compact.

The novel feature of the structure is the manner in which the two chains are held together by the purine and pyrimidine bases. The planes of the bases

are perpendicular to the fibre axis. They are joined together in pairs, a single base from one chain being hydrogen-bonded to a single base from the other chain, so that the two lie side by side with identical z-co-ordinates. One of the pair must be a purine and the other a pyrimidine for bonding to occur. The hydrogen bonds are made as follows: purine position 1 to pyrimidine position 1; purine position 6 to pyrimidine position 6.

If it is assumed that the bases only occur in the structure in the most plausible tautomeric form (that is, with the keto rather than the enol configurations) it is found that only specific pairs of bases can bond together. These pairs are: adenine (purine) with thymine (pyrimidine), and guanine (purine) with cytosine (pyrimidine).

In other words, if an adenine forms one member of a pair, on either chain, then on these assumptions the other member must be thymine; similarly for guanine and cytosine. The sequence of bases on a single chain does not appear to be restricted in any way. However, if only specific pairs of bases can be formed, it follows that if the sequence of bases on one chain is given, then the sequence on the other chain is automatically determined.

It has been found experimentally^{3,4} that the ratio of the amounts of adenine to thymine, and the ratio of guanine to cytosine, are always very close to unity for deoxyribose nucleic acid.

It is probably impossible to build this structure with a ribose sugar in place of the deoxyribose, as the extra oxygen atom would make too close a van der Waals contact.

The previously published X-ray data^{5,6} on deoxyribose nucleic acid are insufficient for a rigorous test of our structure. So far as we can tell, it is roughly compatible with the experimental data, but it must be regarded as unproved until it has been checked against more exact results. Some of these are given in the following communications. We were not aware of the details of the results presented there when we devised our structure, which rests mainly though not entirely on published experimental data and stereochemical arguments.

It has not escaped our notice that the specific pairing we have postulated immediately suggests a possible copying mechanism for the genetic material. Full details of the structure, including the conditions assumed in building it, together with a set of co-ordinates for the atoms, will be published elsewhere.

We are much indebted to Dr. Jerry Donohue for constant advice and criticism, especially on inter-atomic distances. We have also been stimulated by a knowledge of the general nature of the unpublished experimental results and ideas of Dr. M. H. F. Wilkins, Dr. H. R. Franklin and their co-workers at King's College, London. One of us (J.D.W.) has been aided by a fellowship from the National Foundation for Infantile Paralysis.

J. D. WATSON
F. H. C. CRICK

Medical Research Council Unit for the Study of the Molecular Structure of Biological Systems
Cavendish Laboratory, Cambridge.
April 2.

¹ Pauling, L., and Corey, R. B., *Nature*, 237, 348 (1955); *Proc. U.S. Nat. Acad. Sci.*, 38, 54 (1952).

² Furburg, R., *Ann. Chem.*, 586, 434 (1952).

³ Chargaff, E., for references see Zimmoloff, S., *Newsweek*, 6, 402 (1955).


⁴ Wilson, C. B., *J. Am. Physiol.*, 98, 251 (1955).

⁵ Astbury, W. T., *Trans. Faraday Soc.*, 47, 1026 (1951).

⁶ Wilkins, M. H. F., and Randall, J. T., *Nucleic Acids*, 2, 182 (1955).





This figure is a pencil diagram of the DNA double helix. It shows two intertwined chains, one solid and one dashed, representing the sugar-phosphate backbones. Between the chains are horizontal lines representing the base pairs. The bases are on the inside of the helix, and the phosphates are on the outside. The structure is shown as a right-handed helix. A vertical line on the right indicates the fibre axis.






DNA Double Helix



Central Dogma


by F. Crick

DNA → **mRNA** → **Protein ()**



RNA

(Gamow Watson)



■ 20

20

20

■ Geo Gamow ()


“Alpher, Bethe, Gamow”

“ ”


1948 4 1 .

()


DNA



RNA



Protein




Biotechnology & Bioindustry

- **Egypt, Fermentation
(bread, cheese, wine, beer)**
- **1850s, Pure Cultures of Microorganisms**
- **1940s, Random mutation**
- **1973, Recombinant DNA technology**

Recombinant DNA Technology

- Discovery of restriction enzyme and ligase
- Development of recombinant DNA technology
 - Herbert Boyer and Stanley Cohen (1973)



**Cutting and
Joining DNA
Molecules**

DNA




■ 가 :

■ :
DNA

DNA

(1972. 11)



, 1976

■

--

가, 27

■

■ 1976. 4,

■



- 1980. 10. 15


-

- 20 \$35 → \$89

-

가

가



■ &
6,600
가 .



Red BT

**Medical
Biotechnology**



- ()
- t-PA ()
-
- EPO ()
 - -- \$20 / , \$640 가





Green BT

**Agricultural
Biotechnology**



- Polyester가 ()
- ()
- - 1994, Calgene, Flavr-Savr
- Golden Rice (funded by Rockefeller Foundation)
 - A carotene

Bt Designer Crops

- **Bt corn**
- **Bt potato**
- **Bt cotton**
- **Bt soybean**





White BT

**Industrial
Biotechnology**

DuPont – A 203 Year Old Company



E. I. du Pont

