

Electro-Optics:

Propagation of Laser Beams (1)

Yoonchan Jeong

School of Electrical Engineering, Seoul National University

Tel: +82 (0)2 880 1623, Fax: +82 (0)2 873 9953

Email: yoonchan@snu.ac.kr

Wave Equations in Quadratic Index Media (1)

Maxwell's equations:

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0 \quad \nabla \cdot \mathbf{D} = \rho$$

$$\nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = \mathbf{J} \quad \nabla \cdot \mathbf{B} = 0$$

Isotropic, non-magnetic, charge-free, and current-free medium:

$$\nabla^2 \mathbf{E} - \mu \epsilon \frac{\partial^2 \mathbf{E}}{\partial t^2} + (\nabla \log \mu) \times (\nabla \times \mathbf{E}) + \nabla(\mathbf{E} \cdot \nabla \log \epsilon) = 0$$

Wave equation:

$$\nabla^2 \mathbf{E} - \mu \epsilon \frac{\partial^2 \mathbf{E}}{\partial t^2} = -\nabla(\mathbf{E} \cdot \nabla \log \epsilon) \cong 0$$

$$\rightarrow \nabla^2 \mathbf{E} + K^2(\mathbf{r}) \mathbf{E} = 0 \quad \leftarrow K^2(\mathbf{r}) = \omega^2 \mu \epsilon(r)$$

Quadratic index media:

$$K^2(r) = k^2 - k k_2 r^2$$

$$\leftarrow k^2 = K^2(0) = \omega^2 \mu \epsilon(0)$$

Wave Equations in Quadratic Index Media (2)

Scalar-wave approximation:

$$\nabla^2 \mathbf{E} + K^2(r) \mathbf{E} = 0$$

No azimuthal variation:

$$\rightarrow \nabla^2 = \nabla_t^2 + \frac{\partial^2}{\partial z^2} = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{\partial^2}{\partial z^2} \quad \leftarrow \frac{\partial}{\partial \phi} = 0$$

$$\rightarrow E = \psi(x, y, z) e^{-ikz}$$

$$\rightarrow \nabla_t^2 \psi + \cancel{\frac{\partial^2 \psi}{\partial z^2}} - 2ik \frac{\partial \psi}{\partial z} - k^2 \psi + (k^2 - kk_2 r^2) \psi = 0$$

$$\rightarrow \nabla_t^2 \psi - 2ik \psi' - kk_2 r^2 \psi = 0$$

Trial solution:

$$\psi = \exp \left[-i \left(P(z) + \frac{k}{2q(z)} r^2 \right) \right]$$

$$\rightarrow - \left(\frac{k}{q} \right)^2 r^2 - 2i \left(\frac{k}{q} \right)' - k^2 r^2 \left(\frac{1}{q} \right)' - 2kP' - kk_2 r^2 = 0$$

$$\rightarrow \left(\frac{1}{q} \right)^2 + \left(\frac{1}{q} \right)' + \frac{k_2}{k} = 0 \quad \rightarrow \text{For all } r \rightarrow P' = -\frac{i}{q}$$

Gaussian Beams in a Homogeneous Medium (1)

For a homogeneous medium: $\rightarrow k_2 = 0$

$$\rightarrow \left(\frac{1}{q} \right)^2 + \left(\frac{1}{q} \right)' = 0 \quad \rightarrow \quad P' = -\frac{i}{q}$$

$$\rightarrow \frac{1}{q} = \frac{1}{u} \frac{du}{dz} \quad \rightarrow \quad \frac{d^2 u}{dz^2} = 0 \quad \rightarrow \quad u = az + b \quad \left| \begin{array}{l} \rightarrow P' = -\frac{i}{q} = -\frac{i}{z + q_0} \\ \rightarrow P = -i \ln \left(1 + \frac{z}{q_0} \right) \end{array} \right.$$

$$\rightarrow \frac{1}{q} = \frac{a}{az + b} = \frac{1}{z + q_0} \quad \rightarrow \quad q = z + q_0$$

Scalar-wave solution:

$$\psi = \exp \left\{ -i \left[-i \ln \left(1 + \frac{z}{q_0} \right) + \frac{k}{2(z + q_0)} r^2 \right] \right\}$$

$$\rightarrow q_0 \equiv i \frac{\pi \omega_0^2 n}{\lambda} \equiv iz_0 \quad \rightarrow \quad \lambda = \frac{2\pi n}{k}$$

← Choice of a pure imaginary?

Gaussian Beams in a Homogeneous Medium (2)

Scalar-wave solution:

$$\begin{aligned}
 \psi &= \exp\left\{-i\left[-i \ln\left(1 + \frac{z}{q_0}\right) + \frac{k}{2(z+q_0)} r^2\right]\right\} \quad \leftarrow q_0 = iz_0 \\
 &\rightarrow \exp\left[-\ln\left(1 - i\frac{z}{z_0}\right)\right] = \frac{1}{\sqrt{1+(z/z_0)^2}} \exp\left[i \tan^{-1}\left(\frac{z}{z_0}\right)\right] \\
 &\rightarrow \exp\left[\frac{-ikr^2}{2(z+iz_0)}\right] = \exp\left\{\frac{-r^2}{\omega_0^2[1+(z/z_0)^2]} - \frac{ikr^2}{2z[1+(z_0/z)^2]}\right\} \\
 &\rightarrow \omega^2(z) = \omega_0^2\left(1 + \frac{z^2}{z_0^2}\right), \quad R(z) = z\left(1 + \frac{z_0^2}{z^2}\right), \quad \eta(z) = \tan^{-1}\left(\frac{z}{z_0}\right) \\
 &\rightarrow \frac{1}{q(z)} = \frac{1}{z+iz_0} = \frac{1}{R(z)} - i\frac{2}{k\omega^2(z)} = \frac{1}{R(z)} - i\frac{\lambda}{\pi n \omega^2(z)}
 \end{aligned}$$


Gaussian beam solution:

$$\begin{aligned}
 E(x, y, z) &= \psi(x, y, z) e^{-ikz} = E_0 \frac{\omega_0}{\omega(z)} \exp\left\{-i[kz - \eta(z)] - i\frac{kr^2}{2q(z)}\right\} \\
 &= E_0 \frac{\omega_0}{\omega(z)} \exp\left\{-i[kz - \eta(z)] - r^2\left(\frac{1}{\omega^2(z)} + \frac{ik}{2R(z)}\right)\right\}
 \end{aligned}$$

Gaussian Beams in a Homogeneous Medium (3)

Gaussian beam:

$$E(x, y, z) = E_0 \frac{\omega_0}{\omega(z)} \exp \left\{ -i[kz - \eta(z)] - r^2 \left(\frac{1}{\omega^2(z)} + \frac{ik}{2R(z)} \right) \right\}$$



Spherical wave:

$$\begin{aligned} E &\propto \frac{1}{R} e^{-ikR} = \frac{1}{R} \exp \left(-ik\sqrt{x^2 + y^2 + z^2} \right) \\ &\approx \frac{1}{R} \exp \left[-ikz - ik \left(\frac{x^2 + y^2}{2R} \right) \right] \quad \leftarrow x^2 + y^2 \ll z^2 \end{aligned}$$

Energy propagation and angular beam spread:

$$r^2 = x^2 + y^2 = \omega^2(z) \rightarrow r \approx \frac{\lambda}{\pi\omega_0 n} z \rightarrow \text{For large } z$$

$$\rightarrow \theta_{beam} = \tan^{-1} \left(\frac{\lambda}{\pi\omega_0 n} \right) \approx \frac{\lambda}{\pi\omega_0 n} \rightarrow \theta_{beam}\omega_0 \approx \frac{\lambda}{\pi n} = const.$$

Invariant

Fundamental Gaussian Beam in a LL Medium

For a lenslike medium: $\rightarrow k_2 \neq 0$

$$\left(\frac{1}{q}\right)^2 + \left(\frac{1}{q}\right)' + \frac{k_2}{k} = 0, \quad P' = -\frac{i}{q}$$

$$\rightarrow \frac{1}{q} = \frac{u'}{u} \quad \rightarrow u'' + u\left(\frac{k_2}{k}\right) = 0$$

$$\rightarrow u(z) = a \sin \sqrt{\frac{k_2}{k}} z + b \cos \sqrt{\frac{k_2}{k}} z$$

$$\rightarrow u'(z) = a \sqrt{\frac{k_2}{k}} \cos \sqrt{\frac{k_2}{k}} z - b \sqrt{\frac{k_2}{k}} \sin \sqrt{\frac{k_2}{k}} z$$

Complex beam radius:

$$q(z) = \frac{u}{u'} = \frac{q_0 \cos \sqrt{\frac{k_2}{k}} z + \sqrt{k/k_2} \sin \sqrt{\frac{k_2}{k}} z}{-q_0 \sqrt{k_2/k} \sin \sqrt{\frac{k_2}{k}} z + \cos \sqrt{\frac{k_2}{k}} z}$$

$$\rightarrow \frac{1}{q(z)} = \frac{1}{R(z)} - i \frac{\lambda}{\pi n \omega^2(z)}$$

High-Order Gaussian Beam Modes

Scalar-wave approximation:

$$\begin{aligned}\nabla^2 \mathbf{E} + K^2(r) \mathbf{E} &= 0 \\ \rightarrow \nabla^2 &= \nabla_t^2 + \frac{\partial^2}{\partial z^2} = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \\ \rightarrow \nabla_t^2 \psi - 2ik\psi' - kk_2 r^2 \psi &= 0\end{aligned}$$

For a homogeneous medium: $\rightarrow k_2 = 0$

$$\begin{aligned}E_{l,m}(x, y, z) &= E_0 \frac{\omega_0}{\omega(z)} H_l\left(\sqrt{2} \frac{x}{\omega(z)}\right) H_m\left(\sqrt{2} \frac{y}{\omega(z)}\right) \exp\left\{-i \frac{k(x^2 + y^2)}{2q(z)} - ikz + i(l+m+1)\eta\right\} \\ &= E_0 \frac{\omega_0}{\omega(z)} H_l\left(\sqrt{2} \frac{x}{\omega(z)}\right) H_m\left(\sqrt{2} \frac{y}{\omega(z)}\right) \exp\left\{-\frac{x^2 + y^2}{\omega^2(z)} - \frac{ik(x^2 + y^2)}{2R(z)} - ikz + i(l+m+1)\eta\right\}\end{aligned}$$

For a lenslike medium: $\rightarrow k_2 \neq 0$

$$\begin{aligned}E_{l,m}(x, y, z) &= E_0 H_l\left(\sqrt{2} \frac{x}{\omega}\right) H_m\left(\sqrt{2} \frac{y}{\omega}\right) \exp\left(-\frac{x^2 + y^2}{\omega^2}\right) \exp(-i\beta_{l,m}) \\ \rightarrow \beta_{l,m} &= k \left[1 - \frac{2}{k} \sqrt{\frac{n_2}{n_0}} (l+m+1)\right]^{1/2}\end{aligned}$$

H.W.