

Chemical characteristics of water I

Major ionic species in water

Cations

Calcium (Ca^{2+})

Magnesium (Mg^{2+})

Sodium (Na^+)

Potassium (K^+)

Anions

Bicarbonate (HCO_3^-)

Sulfate (SO_4^{2-})

Chloride (Cl^-)

- Derived from contact of water with mineral deposits
- Relatively high in groundwater, low in surface water
- Determining the accuracy of water ion content analysis:

$$\left| \sum \text{anions} - \sum \text{cations} \right| \leq \left(0.1065 + 0.0155 \sum \text{anions} \right)$$

** $\sum$ values in meq/L*

- Most dissolved inorganics are in ionic form
 - Major nonionic: silica (SiO_2)

Dissolved ion analysis

Q: Determine the acceptability of the following water analysis.

Cations	Conc. (mg/L)	Anions	Conc. (mg/L)
Ca ²⁺	93.8	HCO ₃ ⁻	164.7
Mg ²⁺	28.0	SO ₄ ²⁻	134.0
Na ⁺	13.7	Cl ⁻	92.5
K ⁺	30.2		

Major ionic species in water

Firstly, calculate concentrations in meq/L units:

(conc. in meq/L) = (conc. in mg/L) / (Ionic weight, IW) x (oxidation number)

Cations	IW (g/mole)	Conc. in mg/L	Conc. in meq/L
Ca ²⁺	40.1	93.8	4.68
Mg ²⁺	24.3	28.0	2.30
Na ⁺	23.0	13.7	0.60
K ⁺	39.1	30.2	0.77
			Σ(cations) = 8.35

Cations	IW (g/mole)	Conc. in mg/L	Conc. in meq/L
HCO ₃ ⁻	61.0	164.7	2.70
SO ₄ ²⁻	96.1	134.0	2.79
Cl ⁻	35.5	92.5	2.61
			Σ(anions) = 8.10

Major ionic species in water

$$\left| \sum anions - \sum cations \right| \leq \left(0.1065 + 0.0155 \sum anions \right)$$

$$\left| \sum (anions) - \sum (cations) \right| = 0.25$$

$$0.1065 + 0.0155 \sum (anions) = 0.23$$

Therefore,

$$\left| \sum anions - \sum cations \right| > \left(0.1065 + 0.0155 \sum anions \right) \quad (\text{not acceptable})$$

Source of error:

- measurement error of one or more ions
- missing one or more significant ions

Minor ionic species in water

Cations

Aluminum (Al^{3+})	Copper (Cu^{2+})
Ammonium (NH_4^+)	Iron, ferrous (Fe^{2+})
Arsenic (As^+)	Iron, ferric (Fe^{3+})
Barium (Ba^{2+})	Manganese (Mn^{2+})
Borate (BO_4^{3-})	

Anions


Bisulfate (HSO_4^-)	Nitrite (NO_2^-)
Bisulfite (HSO_3^-)	Phosphate, mono- (H_2PO_4^-)
Carbonate (CO_3^{2-})	Phosphate, di- (HPO_4^{2-})
Fluoride (F^-)	Phosphate, tri- (PO_4^{3-})
Hydroxide (OH^-)	Sulfide (S^{2-})
Nitrate (NO_3^-)	Sulfite (SO_3^{2-})

- Mostly derived from contact of the water with mineral deposits
- Some from bacterial and algal activity (ex: NH_4^+ , NO_3^- , NO_2^- , CO_3^{2-} , S^{2-})

Nutrients

- **N & P as major nutrients of interest**
 - Essential for life
 - Most often limiting nutrients in the environment
- **Nitrogen (N)**
 - Exist in various oxidation states: +5, +3, +2, +1, 0, -2, -3
 - Important nitrogen-containing compounds for water quality
 - Organic nitrogen, ammonia (NH_3), nitrite (NO_2^-), nitrate (NO_3^-), urea [$\text{CO}(\text{NH}_2)_2$], nitrogen gas (N_2)

Nitrogen cycle in the environment




Nitrogen cycle

- Uptake by organisms
 - Uptake by microorganisms and plants: NH_3 (most common), NO_3^- , $\text{N}_2 \rightarrow$ produce proteins
 - Conversion of N_2 to organic-N by bacteria is called “nitrogen fixation” (by limited number of bacterial species)
 - Human contribution to nitrogen cycle: Haber-Bosch process
$$\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$$
 - Uptake by animals and humans: nitrogen must be in organic form (protein)
- Release from organisms
 - Animals excrete urea and other forms of organic-N (ex: proteins)
 - Dead organisms $\rightarrow$ release organic-N into the environment


Nitrogen cycle

- Fate of N released into the environment
 - Organic-N is degraded by bacteria to urea and NH_3
 - Urea is easily hydrolyzed to NH_3

Urea hydrolysis




- Ammonia is oxidized serially by certain groups of bacteria:



<nitrification>

Nitrogen cycle

- Nitrate and nitrite is reduced by various types of bacteria to produce nitrogen gas (N_2) by series of reactions:



<denitrification>


- Note nitrous oxide (N_2O) is a potent greenhouse gas (greenhouse gas potential 265-298 time greater than CO_2)
- N_2O may be released as an intermediate of both nitrification and denitrification

Measurement of N in water

- Each ionic species can be measured by ion chromatography or colorimetric methods
- Organic nitrogen is determined by the Kjeldahl method: organic-N is degraded by acid and heat to ammonium and then ammonium content is determined
- **Total Kjeldahl nitrogen (TKN)** = organic-N + ammonia-N
- To determine organic-N only by the Kjeldahl method, the water is first heated to remove NH_3 by volatilization


General methods for measuring ions

#2



Colorimetric method

- Add chemical agents that will react with the compound to be measured to form products that have a color
- Measure absorbance by spectrophotometer or compare the color with standards



IC at Water Quality & Environment Lab., SNU

Ion chromatography (IC)

- Sample is injected to a column which has different affinity to different ions
- An eluent continuously flushes the column and the ions flow out of the column at different times
- Concentration of each ion is determined by measuring electrical conductivity

Phosphorus (P)

- Used..
 - in fertilizers
 - for corrosion control in water supply and industrial cooling water
 - in synthetic detergents
- P-containing compounds relevant to water quality
 - Orthophosphates: PO_4^{3-} , HPO_4^{2-} , H_2PO_4^- , H_3PO_4
 - Can be directly utilized by organisms
 - Easily measured by colorimetric methods / ion chromatography
 - Polyphosphates ($(\text{PO}_3)_6^{3-}$, $\text{P}_3\text{O}_{10}^{5-}$, $\text{P}_2\text{O}_7^{4-}$, ...) and organic phosphates
 - Needs breakdown to orthophosphates for biological metabolism / analysis


Sulfur (S)

- Essential element for life

– C, H, O, N, **S**, P, K, ...



– Required in the synthesis of proteins, released when protein degrades

- Reduced biologically under anaerobic conditions



– Anaerobic conditions occur in sediment, subsurface, sewers, and anaerobic processes in wastewater treatment

– The sulfide ion (S^{2-}) may combine with hydrogen to form hydrogen sulfide gas (H_2S)



pH

$$pH = -\log_{10}[H^+]$$

- Ionization constant of water

$$[H^+][OH^-] = K_w \quad K_w = 10^{-14} \text{ at } 25^\circ\text{C}$$

$$p \equiv -\log_{10} \rightarrow pH + pOH = 14 \text{ at } 25^\circ\text{C}$$

Q: pH in pure H₂O at 25°C?

Electrical conductivity (EC)

- A measure of an ability of a solution to conduct an electrical current
- Unit: millisiemens per meter (mS/m) or microsiemens per centimeter ($\mu\text{S}/\text{cm}$)
- Electrical current is transported by ions in a solution $\rightarrow$ related to the concentration of ions in a solution



Conductivity meter & probe

Electrical conductivity (EC)

- Conversion between EC and ionic concentration
 - Conc. of each ionic species in water and EC

$$EC \cong \sum_i (C_i \times f_i)$$

EC = electrical conductivity ($\mu\text{S}/\text{cm}$)

C_i = conc. of ionic species i in solution (meq/L)

f_i = conversion factor

Cations	f_i [$(\mu\text{S}/\text{cm}) \cdot (\text{meq/L})^{-1}$]	Anions	f_i [$(\mu\text{S}/\text{cm}) \cdot (\text{meq/L})^{-1}$]
Ca^{2+}	52.0	HCO_3^-	43.6
Mg^{2+}	46.6	CO_3^{2-}	84.6
K^+	72.0	Cl^-	75.9
Na^+	48.9	NO_3^-	71.0
		SO_4^{2-}	73.9

Electrical conductivity (EC)

- Conversion between EC and ionic concentration
 - Applying generic composition of ionic species in water, EC can be used to estimate the ionic strength and TDS of a solution

$$I = EC \text{ (in } \mu\text{S/cm)} \times (1.6 \times 10^{-5})$$

Tchobanoglous & Schroeder (1985) Water Quality

$$TDS \text{ (mg/L)} = EC \text{ (in } \mu\text{S/cm)} \times (0.55 - 0.70)$$

Metcalf, Eddy, AECOM (2014) Wastewater Engineering

Alkalinity

- The capacity of water to neutralize acid
- Determined by titrating water with a strong acid to pH=4.5

$$\begin{aligned} \text{Alk (eq/L)} &= (\text{HCO}_3^-) + (\text{CO}_3^{2-}) + \cdots + (\text{OH}^-) - (\text{H}^+) \\ &= [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + \cdots + [\text{OH}^-] - [\text{H}^+] \end{aligned}$$

Include B(OH)_4^- ,
 PO_4^{3-} , HPO_4^{2-} ,
 SiO(OH)_3^- , etc. if
significant

- Most of the time, practically:

$$\text{Alk (eq/L)} \cong [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{OH}^-]$$

- Most of the time, at neutral pH:

$$\text{Alk (eq/L)} \cong [\text{HCO}_3^-]$$


More common unit for Alk:
“mg/L as CaCO_3 ”

Conversion

$\text{Alk (in mg/L as CaCO}_3\text{)}$
 $= \text{Alk (in meq/L)} \times 50 \text{ mg CaCO}_3/\text{meq}$

Hardness

- The term used to characterize a water that does not lather well (react with soap to form a scum)
- Caused by polyvalent cations in water (+2, +3, ...); mostly Ca^{2+} & Mg^{2+}
- These ions are also easily precipitated to produce scales in pipes transporting hot water




#5



#6

CaCO₃ precipitation – temperature effect

Recall:



K_{a2} increases with increasing temperature:

$$K_{a2} = 2.75 \times 10^{-11} \quad (\text{at } 5^\circ\text{C})$$

$$K_{a2} = 6.03 \times 10^{-11} \quad (\text{at } 40^\circ\text{C})$$

Also recall:

$$K_{a2} = \frac{[CO_3^{2-}][H^+]}{[HCO_3^-]}, \quad [CO_3^{2-}] = K_{a2} \frac{[H^+]}{[HCO_3^-]}$$

➡ Higher CO₃⁻² fraction when water is heated,
Ca²⁺ is more likely to be precipitated as CaCO₃

Hardness

- Total hardness (TH)

- Technically: the sum of all polyvalent cations

$$TH(eq/L) = (Ca^{2+}) + (Mg^{2+}) + (Fe^{3+}) + (Fe^{2+}) + (Ba^{2+}) + \dots = \sum_{i=1}^n (X^{m+})_i$$

- Practically (most of the time): sum of Ca^{2+} & Mg^{2+}

$$TH(eq/L) \cong (Ca^{2+}) + (Mg^{2+}) = 2[Ca^{2+}] + 2[Mg^{2+}]$$


“mg/L as $CaCO_3$ ” is more common for hardness as well!

- Carbonate hardness (CH) and noncarbonate hardness (NCH)

- CH: the maximum amount of hardness that can be associated with carbonates (HCO_3^- and CO_3^{2-})
- $NCH = TH - CH$
- When **TH > Alk**: **CH = Alk**, **NCH = TH - CH**
- When **TH ≤ Alk**: **CH = TH**, **NCH = 0**

Why are we interested in CaCO_3 ?

Recall for the following precipitation reaction:



Inspect the pK_{sp} of potential Ca/Mg precipitates:

$$pK_{sp} (\text{CaCO}_3) = 8.55$$

$$pK_{sp} (\text{MgCO}_3) = 7.46$$

$$pK_{sp} (\text{Ca(OH)}_2) = 5.26$$

$$pK_{sp} (\text{Mg(OH)}_2) = 10.74$$




Sodium adsorption ratio (SAR)

- Related to the agricultural production
 - Important property for irrigation water
- High sodium (Na^+) content in soil reduces soil permeability!
 - Most clay surfaces are negatively (-) charged
 - Cations are attached to clay surfaces
 - Attachment of Na^+ ion on clay surfaces
 - swelling of clay by introduction of water molecules between clay sheets
 - soil pore size ↓
 - soil permeability ↓
 - crop productivity ↓
 - So, irrigation of water with high Na^+ content can result in replacement of Ca^{2+} and Mg^{2+} in soil, resulting in low crop productivity

Clay swelling by water addition



#7



Sodium adsorption ratio (SAR)

$$SAR = \frac{(Na^+)}{\sqrt{\frac{(Ca^{2+}) + (Mg^{2+})}{2}}}$$

Note:

Here, () denotes meq/L,
not eq/L

SAR < 3: low risk

$3 \leq SAR \leq 6$: slight to moderate risk

SAR > 6: high risk

References

- #1) *Davies, M. L., Masten, S. J. (2014) Principles of Environmental Engineering and Science, 3rd ed. McGraw-Hill (figure provided by the publisher).*
- #2) <http://stuff.iorodeo.com/docs/ammonia/api.html>
- #3) <https://derangedphysiology.com/main/cicm-primary-exam/required-reading/acid-base-physiology/acid-base-disturbances/Chapter%20613/sulfate-anion-its-origins-and-its-clearance>
- #4) <https://www.coleparmer.co.uk/p/oakton-pc-700-benchtph-conductivity-meters/58971>
- #5) <https://blogs.ubc.ca/communicatingscience2018w112/2018/11/12/is-hard-water-bad-for-your-health/>
- #6) <https://laundryledger.com/how-hard-water-affects-the-laundry-process/>
- #7) *David, S. (2005) The Effects of High Salinity Groundwater on the Performance of Clay Barriers. SKI Report 2005:54, p. 5.*