

Chemical characteristics of water II

Organic content

- Contaminated water contains various kinds of organic compounds
 - Proteins, carbohydrates, fats and oils, urea, etc. from food and human wastes
 - Synthetic organic compounds
 - Organics released to waters → consumption of dissolved oxygen by microorganisms → anaerobic (septic) condition → destroy aquatic environment (ex: fish kills), odor problems, production of toxic compounds, etc.
 - Removal of organic compounds is one of the major target for wastewater treatment
- Measurement of organic content as a whole
 - Biochemical oxygen demand (BOD)
 - Chemical oxygen demand (COD)
 - Total organic carbon (TOC)

BOD

- Measurement of dissolved oxygen used by microorganisms in the biochemical oxidation of organic matter
- **BOD test procedure**
 - The water sample is diluted such that the difference between the DO before and after the test can be determined (estimated BOD: 2-6 mg/L)
 - The diluted water sample is inoculated with microorganisms that degrade organic matter
 - The diluted, inoculated water sample is incubated for a certain time period (usually 5 days)
 - The DO before and after the incubation is measured to determine the BOD of the sample



BOD


- Modeling BOD reaction: assume first-order reaction

$$\frac{d(L_t)}{dt} = -k_1 L_t$$

L_t = amount of organics remaining at time t (d) expressed in oxygen equivalents (mg O_2 /L)
 k_1 = first-order rate constant (1/d)

Integrating from $t=0$ to t ,

$$L_t = L_0(e^{-k_1 t})$$



Note

$$BOD_t = L_0 - L_t$$

$$L_0 = UBOD$$

$UBOD$ = ultimate BOD (mg/L)

BOD

$$\rightarrow BOD_t = UBOD - L_t = UBOD(1 - e^{-k_1 t})$$

BOD_t = the BOD value at time t (mg/L)

- Temperature effect
 - modified van't Hoff-Arrhenius relationship:

$$k_{1T} = k_{1_{20}} \theta^{T-20}$$

T = temperature in °C

- Typically used value of θ :
1.056 (20-30°C) / 1.135 (4-20°C)

NBOD vs. CBOD

- Ammonia-nitrogen in wastewater may significantly contribute to the total oxygen demand by nitrification:





Figure 2-24

Definition sketch for the exertion of the carbonaceous and nitrogenous biochemical oxygen demand in a waste sample.



Metcalf, Eddy, AECOM (2014)

NBOD vs. CBOD

- The oxygen demand associated with the oxidation of ammonia is referred to as nitrogenous biochemical oxygen demand (NBOD)
- Carbonaceous biochemical oxygen demand (CBOD): the oxygen demand associated with the oxidizable carbon in the sample
- When NBOD is significant, nitrification is suppressed by adding chemical agents for the measurement of CBOD


COD

- Measured by oxidizing the organic compounds in water using a strong oxidizing agent
- Oxidizing agent: potassium dichromate (K_2CrO_7 - more common) or potassium permanganate ($KMnO_4$)
- Can be fractionated into particulate and soluble COD
 - Soluble COD: readily biodegradable / nonbiodegradable
 - Particulate COD: slowly biodegradable / nonbiodegradable



COD

- **COD > BOD** because:
 - Many organics that are difficult to be oxidized biologically can be oxidized chemically (ex: lignin)
 - Inorganic substances in water may be oxidized by chemical oxidizing agents
 - Certain organic substances may be toxic to microorganisms used in the BOD test
 - When microorganisms grow, they utilize some fraction of organic compounds to synthesize cells instead of oxidizing them



TOC



- Measures all organic carbon in a water sample including those that cannot be chemically/biologically oxidized
- Can be fractionated into particulate/soluble TOC
- Three steps for measurement
 - **Acidification:** add acid to reduce the pH → removes carbonate species (inorganic carbon) from water
 - **Oxidation:** use heat, oxygen, ultraviolet radiation, or combination of those to oxidize organic carbon to CO₂
 - **Quantification:** measure the amount of CO₂ production with an infrared analyzer or other means
- TOC: measures amount of C / BOD & COD: measures amount of O₂ consumed by oxidation
 - different COD/TOC ratio for different compounds!

BOD, COD, & TOC

Q: Determine the theoretical ratios of BOD_5/COD and COD/TOC for an organic compound represented by $C_5H_7O_2N$. Use the following assumptions:

- The compound can be completely mineralized biologically
- Only CBOD is considered for BOD
- The BOD first-order reaction rate constant, k_1 , is 0.23/d

Individual organic compounds

- Some organic compounds have particular toxicity to humans and aquatic organisms → have to be regulated individually
- Sources
 - Commercial and industrial wastewater
 - Disinfection byproducts
 - Surface runoff from agricultural land (ex: pesticides)
 - Surface runoff from urban area (ex: oil spill, additives used for vehicles, sealant for pavements)
 - Pharmaceuticals and personal care products (PPCPs)
 - Mostly not regulated currently, but of recent interest

