

# Redox reactions


# Oxidation/reduction reactions

---

- **Electron transfer reactions**

- Redox couples & half reactions

- One component is reduced (accepts electron(s))
- One component is oxidized (donates electron(s))



- Most natural organic compounds are (primarily) electron donors

# Oxidation/reduction reactions




---

- **Most important biotic reactions**

- Primary energy source

- Cell maintenance
- Cell growth

ex) glucose oxidation mediated by microorganisms




# Oxidation/reduction reactions

---

- **Photochemical reactions**
  - Will discuss later
- **Redox reactions also occur abiotically in the dark**
  - May involve mediators/catalysts
  - Although the reaction itself is abiotic, biological molecules can be involved

# Oxidation/reduction reactions

- Reduction of organic pollutants mediated by biological molecules




# Oxidation/reduction reactions

---

- Pd-catalyzed hydrodehalogenation

## Net Reaction:




- Rapid reduction of TCE at the surface of Pd



# Thermodynamics of redox reactions

---


- Consider a reversible reactions to convert 1,4-benzoquinone (BQ) to hydroquinone (HQ):



- Use reduction potentials for evaluating the free energy of the half reaction
  - Perform the reaction at the surface of an inert electrode (ex: platinum, graphite)
  - At the other side, another inert electrode is immersed in an aqueous solution maintained at pH 0 (i.e.,  $\{\text{H}^+\} = 1$ ) and bubbled with molecular hydrogen ( $p_{\text{H}_2} = 1 \text{ bar}$ ) – standard hydrogen electrode (SHE)

# Thermodynamics of redox reactions

---







# Thermodynamics of redox reactions

---

- **Overall reaction:**



- **With half reactions:**



# Thermodynamics of redox reactions

---

- **Relationship between the electrical potential and the free energy**

- Assuming electrochemical equilibrium at the electrode surface, then the potential difference,  $\Delta E$ , is directly related to the free energy change,  $\Delta_r G$  of the reaction:

$$\Delta_r G = -nF\Delta E$$


$n$  = number of electrons transferred

$F$  = Faraday constant, 96485 Coulomb/mole

= 96.5 kJ/mole-V

- **At the SHE**

- We assign a zero value of electrical potential, and thus, a zero value of a standard free energy change:



- Then, the electrical potential change of the half reaction,  $BQ + 2H^+ + 2e^- \rightleftharpoons HQ$ , can be directly measured by the electrical potential change at the galvanic cell

# Thermodynamics of redox reactions

---

- The electrical potential relative to SHE,  $E_H$

$$\Delta_r G = -nFE_H$$

- The  $E_H$  value for any conditions other than the “standard” state

$$E_H = E_H^0 - \frac{RT}{nF} \ln Q_r = E_H^0 - \frac{2.303RT}{nF} \log Q_r$$

- $E_H^0$  = standard redox potential or standard reduction potential ( $E_H$  at 25°C with unit (1) activities for all reaction components)
- $Q_r$  = reaction quotient

# Conditions for $E_H^0$ & $E_H^0 (W)$


---

- $E_H^0$ 
  - All reaction components have unit activities
- $E_H^0 (W)$ 
  - $E_H$  under **typical natural water conditions**:
    - pH = 7
    - $[Cl^-] = 10^{-3}$  M;  $[Br^-] = 10^{-5}$  M
    - Organic oxidant and reductant have unit activities

# $\Delta_r G^0$ & $\Delta_r G^0 (W)$ and $E_H^0$ & $E_H^0 (W)$

---

**Q:** Consider the half reaction in aqueous solution (denitrification):



Calculate the  $E_H^0$ ,  $E_H^0 (W)$ , and  $\Delta_r G^0 (W)$  values of the reaction. Initiate with the calculation of  $\Delta_r G^0$  using the enthalpy of formation for each species:

$$\Delta_f G^0_{NO_3^-} (aq) = -111.3 \text{ kJ/mole}$$

$$\Delta_f G^0_{H^+} (aq) = 0 \text{ kJ/mole}$$

$$\Delta_f G^0_{e^-} (aq) = 0 \text{ kJ/mole}$$

$$\Delta_f G^0_{N_2} (g) = 0 \text{ kJ/mole}$$

$$\Delta_f G^0_{H_2O} (l) = -237.2 \text{ kJ/mole}$$

# Oxidation/reduction reactions

**Table 14.2** Standard Reduction Potentials and Average Standard Free Energies of Reaction (per Electron Transferred) at 25 °C of Some Redox Couples that Are Important in Natural Redox Processes (The reactions are ordered in decreasing  $E_H^0(W)$  values.)<sup>a</sup>

Halfreaction		$E_H^0$ (V)	Water @ pH=7 <sup>-</sup>	
Oxidized Species	Reduced Species		$E_H^0(W)$ (V)	$\Delta_r G^0(W)/n^c$ (kJ·mol <sup>-1</sup> )
(1a)	$O_2(g) + 4 H^+ + 4 e^- = 2 H_2O$	+1.23	+0.81	-78.3
(1b)	$O_2(aq) + 4 H^+ + 4 e^- = 2 H_2O$	+1.19	+0.77	-74.3
(2)	$2 NO_3^- + 12 H^+ + 10 e^- = N_2(g) + 6 H_2O$	+1.24	+0.74	-72.1
(3)	$MnO_2(s) + HCO_3^- (10^{-3}) + 3 H^+ + 2 e^- = MnCO_3(s) + 2 H_2O$		+0.53 <sup>b</sup>	-50.7 <sup>b</sup>
(4)	$NO_3^- + 2 H^+ + 2 e^- = NO_2^- + H_2O$	+0.85	+0.43	-41.6
(5)	$NO_3^- + 10 H^+ + 8 e^- = NH_4^+ + 3 H_2O$	+0.88	+0.36	-35.0
(6)	$FeOOH(s) + HCO_3^- (10^{-3} M) + 2 H^+ + e^- = FeCO_3(s) + 2 H_2O$		-0.05 <sup>b</sup>	+ 4.8 <sup>b</sup>
(7)	$CH_3COCOO^-$ (pyruvate) + 2 H <sup>+</sup> + 2 e <sup>-</sup> = CH <sub>3</sub> CHOHCOO <sup>-</sup> (lactate)		-0.19	+17.8
(8a)	$HCO_3^- + 9 H^+ + 8 e^- = CH_4(aq) + 3 H_2O$	+0.21	-0.20	+19.3
(8b)	$CO_2(g) + 8 H^+ + 8 e^- = CH_4(g) + 2 H_2O$	+0.17	-0.24	+23.6
(9)	$SO_4^{2-} + 9 H^+ + 8 e^- = HS^- + 4 H_2O$	+0.25	-0.22	+20.9
(10)	$S(s) + 2 H^+ + 2 e^- = H_2S(aq)$	+0.14	-0.27	+26.0
(11a)	$2 H^+ + 2 e^- = H_2(aq)$	+0.08	-0.33	+31.8
(11b)	$2 H^+ + 2 e^- = H_2(g)$	0.00	-0.41	+40.0
(12)	$6 CO_2(g) + 24 H^+ + 24 e^- = C_6H_{12}O_6(\text{glucose}) + 6 H_2O$	-0.01	-0.43	+41.0

<sup>a</sup> Note that most of the electron transfer reactions involving these redox couples are biologically mediated.


Data from Thauer et al. (1977) and Stumm and Morgan (1995). <sup>b</sup> Note that these values correspond to  $[HCO_3^-] = 10^{-3} M$ .

<sup>c</sup>  $n$  = number of electrons transferred.

# Thermodynamics of redox reactions

---

- **The overall free energy change of a redox reaction:**



$$\Delta_rG^0 = \Delta_{red}G^0 + \Delta_{oxi}G^0$$

- Under the standard conditions:
  - Negative  $\Delta_rG^0$ : favorable reaction
  - Large  $|\Delta_rG^0|$  (*with*  $\Delta_rG^0 < 0$ ): strong driving force for the forward reaction, makes the reversible reaction difficult
  - Note: this is thermodynamics, **NOT** kinetics!

# A simple redox reaction

---

	$\Delta G^0(W)/e^-$ , kJ/mole
<b>A</b> $2H^+ + 2e^- \leftrightarrow H_2(g)$	+40.0
<b>B</b> $O_2(g) + 4H^+ + 4e^- \leftrightarrow 2H_2O$	-78.3
<b>B-2A</b> $2H_2(g) + O_2(g) \leftrightarrow 2H_2O$	

$$\begin{aligned}\Delta_r G^0 &= 4 \times (-78.3 - 40.0) \\ &= -473.2 \text{ kJ/mole } O_2\end{aligned}$$

$$\Delta_r G^0 = -RT \ln(K_{eq})$$

$$K_{eq} = \frac{1}{P_{O_2} P_{H_2}^2} = e^{-(\Delta G_r^0/RT)} = e^{191} = 10^{83}$$



# Oxidation cascade









Figure 14.3 Variation in concentrations of important dissolved redox species along the flowpath of a contaminant plume in groundwater. This sequence results in several zones of characteristic microbial metabolism and corresponding redox conditions (adapted from Bouwer et al., 1984).

*Mineral oxidation/reduction*

# Oxidation cascade – why?

---



# Oxidation cascade – why?

---

<i>Electron Acceptor</i>	<i><math>\Delta G</math>/mole glucose oxidized, kJ</i>
$O_2 \rightarrow H_2O$	-2863.2
$NO_3^- \rightarrow N_2$	-2714.4
$Fe^{3+} \rightarrow Fe^{2+}$	-868.8
$SO_4^{2-} \rightarrow S^{2-}$	-482.4
<i>Methanogenesis</i>	-417.6

*The electron acceptor that generates the most energy get used up first*

*More energy/unit substrate means that more of the substrate can be used to make new cells*

Note: the system analyzed here is pH = 7 with  $[HCO_3^-] = 10^{-3}$  M


# Oxygen cascade: implications

---

- **Different electron acceptor – different microbial community**
  - Different enzymes
  - Different pathways
- **Compound fate may depend on microbial community**
  - Different transformation rates
  - Different products

# Redox reactions – multiple pathways


- **Aerobic microbial transformation of toluene**



Five different aerobic biodegradation pathways for toluene, each initiated through the activity of a mono- or dioxygenase together with molecular oxygen. (Mikesell et al., 1993)

# Redox reactions – multiple pathways


- Anaerobic PCE reaction pathways with zero valent iron ( $\text{Fe}^0$ )



*Hypothesized reaction sequence for reduction of chlorinated ethenes and related compounds by  $\text{Fe}^0$ . Adapted from Arnold and Roberts (2000).*

# Redox reactions – multiple pathways

- Methanogenic TCA transformations




Adapted from  
Vogel et al. (1987)

# Halogenated aliphatic hydrocarbons


---

- **Heavily used:**
  - As solvents
  - As pesticides
  - For fire control
- **Halogens affect:**
  - Size
  - C oxidation state


BROMOETHANE




DBCP




1,2-DICHLOROETHANE




CFC-113



TRICHLOROETHYLENE





1,1,1-TRICHLOROETHANE





# Halogenated aliphatic hydrocarbons













**Table 14.3** Standard Reduction Potentials and Average Standard Free Energies of Reaction (per Electron Transferred) at 25°C of Some Organic Redox Couples in Aqueous Solution (The reactions are ordered in decreasing  $E_{\text{H}}(\text{W})$  values.)<sup>a</sup>

Halfreaction		$E_{\text{H}}^0$ (V)	$E_{\text{H}}^0(\text{W})^b$ (V)	$\Delta_r G^0(\text{W})/n^c$ (kJ·mol <sup>-1</sup> )
Oxidized Species	Reduced Species			
(1) $\text{CCl}_3\text{—CCl}_3 + 2\text{e}^-$	= $\text{Cl}_2\text{C=CCl}_2 + 2\text{Cl}^-$	+ 0.95	+ 1.13	- 109.0
(2) $\text{CBr}_4 + \text{H}^+ + 2\text{e}^-$	= $\text{CHBr}_3 + \text{Br}^-$	+ 0.89	+ 0.83	- 80.1
(3) $\text{CCl}_4 + \text{H}^+ + 2\text{e}^-$	= $\text{CHCl}_3 + \text{Cl}^-$	+ 0.79	+ 0.67	- 64.7
(4) $\text{CHBr}_3 + \text{H}^+ + 2\text{e}^-$	= $\text{CH}_2\text{Br}_2 + \text{Br}^-$	+ 0.67	+ 0.61	- 58.9
(5) $\text{Cl}_2\text{C=CCl}_2 + \text{H}^+ + 2\text{e}^-$	= $\text{Cl}_2\text{C=CHCl} + \text{Cl}^-$	+ 0.70	+ 0.58	- 56.0
(6) $\text{CHCl}_3 + \text{H}^+ + 2\text{e}^-$	= $\text{CH}_2\text{Cl}_2 + \text{Cl}^-$	+ 0.68	+ 0.56	- 54.0
(7)  + $\text{H}^+ + 2\text{e}^-$	=  + $\text{Cl}^-$	+ 0.68	+ 0.56	- 54.0

<sup>a</sup> Estimated from thermodynamic data Dean (1985); Vogel et al. (1987); Krop et al. (1994); Roberts et al. (1996); Totten and Roberts (2001). <sup>b</sup>  $[\text{H}^+] = 10^{-7}$ ,  $[\text{Cl}^-] = 10^{-3}$ ,  $[\text{Br}^-] = 10^{-5}$ . <sup>c</sup>  $n$  = number of electrons transferred.

# Halogenated aliphatic hydrocarbons

**Table 14.3** Standard Reduction Potentials and Average Standard Free Energies of Reaction (per Electron Transferred) at 25°C of Some Organic Redox Couples in Aqueous Solution (The reactions are ordered in decreasing  $E_H^0(W)$  values.)<sup>a</sup>


Halfreaction		$E_H^0$ (V)	$E_H^0(W)^b$ (V)	$\Delta_r G^0(W)/n^c$ (kJ·mol <sup>-1</sup> )
Oxidized Species	Reduced Species			
(8)  + H <sup>+</sup> + 2e <sup>-</sup>	=  + Cl <sup>-</sup>	+ 0.54	+ 0.42	- 40.5
(9)  + 6H <sup>+</sup> + 6e <sup>-</sup>	=  + 2H <sub>2</sub> O	+ 0.83	+ 0.42	- 40.5
(10)  + 2H <sup>+</sup> + 2e <sup>-</sup>	= 	+ 0.70	+ 0.28	- 27.0
(11)  + 2H <sup>+</sup> + 2e <sup>-</sup>	=  + H <sub>2</sub> O	+ 0.57	+ 0.16	- 15.4
(12)  + 4H <sup>+</sup> + 4e <sup>-</sup>	= 2 	+ 0.31	- 0.10	+ 9.7
(13)  + 2H <sup>+</sup> + 2e <sup>-</sup>	=  + H <sub>2</sub> O	+ 0.17	- 0.24	+ 23.2
(14) R-S-S-R + 2H <sup>+</sup> + 2e <sup>-</sup> (cystine)	= 2R-SH (cysteine)	+ 0.02	- 0.39	+ 37.6

<sup>a</sup> Estimated from thermodynamic data Dean (1985); Vogel et al. (1987); Krop et al. (1994); Roberts et al. (1996); Totten and Roberts (2001). <sup>b</sup> [H<sup>+</sup>] = 10<sup>-7</sup>, [Cl<sup>-</sup>] = 10<sup>-3</sup>, [Br<sup>-</sup>] = 10<sup>-5</sup>. <sup>c</sup>  $n$  = number of electrons transferred.


# Halogenated aliphatic hydrocarbons

---


## ex) Reductive dechlorination of $C_2Cl_6$



*Couple with Eq. 11b of Table 14.2:*



*Overall reaction:*



$$\Delta_r G^0 = -183.4 - 0 = -183.4 \text{ kJ/mole } C_2Cl_6$$

# Halogenated aliphatic hydrocarbons

---

## ex) Reductive dechlorination of $C_2Cl_6$

$$\Delta_r G^0 = -RT \ln K_{eq}$$

$$K_{eq} = e^{-\Delta_r G^0 / RT} = 1.4 \times 10^{32} = \frac{[C_2Cl_4][H^+]^2[Cl^-]^2}{[C_2Cl_6]P_{H_2}}$$

Using  $[H^+] = 10^{-7}$  M &  $[Cl^-] = 10^{-3}$  M:

$$\frac{[C_2Cl_4]}{[C_2Cl_6]P_{H_2}} = 1.4 \times 10^{52}$$

*For any reasonable partial pressure of  $H_2$ , the amount of  $C_2Cl_6$  remaining at equilibrium is infinitesimal. Thermodynamically the reaction goes all the way to the right.*