

Organic chemistry background V:  
Hydrogen donors/acceptors,  
EDGs/EWGs

# Hydrogen donors and acceptors


---

- **Hydrogen (bond) donors**

An ion or molecule which possesses a hydrogen atom attached to a relatively electronegative atom such that the hydrogen can participate in a hydrogen bond

- **Hydrogen (bond) acceptors**


An electronegative ion or molecule which possesses a lone electron pair in order to form a hydrogen bond



# Electron donating and withdrawing groups

- A functional group attached to a carbon atom in an organic molecule may affect the reactivity of the molecule

ex) Nitration of a substituted benzene



R	Reaction rate relative to benzene	Product ratio			Comments
		ortho	meta	para	
CH <sub>3</sub>	25	63%	3%	34%	<b>Activated</b> (ortho/para directed)
CF <sub>3</sub>	2.5 x 10 <sup>-5</sup>	6%	91%	3%	<b>Deactivated</b> (meta directed)

# Electron donating groups


---

- Increase the electron density of the aromatic ring
- Make the molecule more nucleophilic (activated)
- The molecule tends to react with electrophiles at ortho- & para-sites

# Electron withdrawing groups

---

- Decrease the electron density of the aromatic ring
- Make the molecule less nucleophilic (deactivated)
- The molecule tends to react with electrophiles at meta-sites

		<i>Most Activating</i>			#1
		$-\ddot{\text{O}}^-$			
		$-\ddot{\text{N}}\text{R}_2$			
		$-\ddot{\text{N}}\text{H}_2$	Strongly Activating		
		$-\ddot{\text{O}}\text{H}$			
		$-\ddot{\text{O}}\text{R}$			
Activating		$-\ddot{\text{N}}\text{HCO}$			
EWG		$-\ddot{\text{O}}\text{CO}$	Moderately Activating	ortho / para directing	
		$-\text{R}$			
			Weakly Activating		
		$-\text{C}=\text{CR}_2$   H			
Reference		$-\text{H}$			
		$-\text{X}$	Weakly Deactivating	=	
		$-\text{CHO}$			
		$-\text{COR}$			
		$-\text{COR}$	Moderately Deactivating		
		$-\text{COOH}$			
		$-\text{COCl}$			
Deactivating		$-\text{CH}_3$		meta directing	
		$\text{C}\equiv\text{N}$			
		$-\text{SO}_2\text{CH}_3$			
		$-\text{SO}_2\text{NH}_2$	Strongly Deactivating		
		$-\text{NH}_3^+$			
		$-\text{NR}_3^+$			
		$-\text{N}^+\text{O}^-$			
		<i>Most Deactivating</i>			

# References

---

#1) <http://www.chem.ucalgary.ca/courses/350/Carey5th/Ch12/ch12-8b.html>

Organic chemistry background VI:  
**Organics with heteroatoms**




# Oxygen-containing functional groups


- Alcohols, phenols and ethers

- Alcohols: R-OH  
(R: alkyl group)
- Phenols: R-OH  
(R: aromatic group)
- Ethers: R<sub>1</sub>-O-R<sub>2</sub>

alcohols (R—OH) and phenols (Ar—OH)



ethers (R<sub>1</sub>—O—R<sub>2</sub>)



# Oxygen-containing functional groups


---

- **Alcohols, phenols and ethers**
  - Oxygen atoms participate in hydrogen bonds: significant changes in physicochemical properties of the molecule
  - R-OH: may act as both H-donor and H-acceptor
  - R<sub>1</sub>-O-R<sub>2</sub>: acts only as an H-acceptor
  - Dissociation of a R-OH group
    - R-OH group may dissociate in water (renders H<sup>+</sup>) → act as a weak acid
    - Especially for phenols
    - Greater dissociation tendency for phenols substituted with electron-withdrawing substituents

# Dissociation of phenols

---

Compound	pK <sub>a</sub>	Dominant species at pH=7.0
phenol	9.95	protonated (>99.9%)
2,4-dichlorophenol	7.90	protonated (~89%)
pentachlorophenol	4.90	deprotonated (>99.9%)



# Oxygen-containing functional groups

---

- Aldehyde and keto groups

- C=O bonds
- Aldehyde: C-CHO; keto: R<sub>1</sub>-CO-R<sub>2</sub>
- H-acceptors
- Quite reactive


#2




formaldehyde  
(disinfectant,  
chemical intermediate)




acetaldehyde  
(chemical intermediate,  
solvent)



isobutyraldehyde  
(chemical intermediate,  
solvent, disinfection  
byproduct in drinking water)




benzaldehyde  
(chemical intermediate,  
solvent)




acrolein  
(chemical intermediate  
for polymer production)



acetone  
(chemical intermediate,  
solvent)



2-butanone  
(solvent)





methyl phenylketone  
(acetophenone; chemical  
intermediate, solvent)


# Oxygen-containing functional groups

- **Carboxylic groups**


- R-COOH
- May dissociate in aqueous solution ( $pK_a$  in the range of 0-6)
- Both strong H-donors and acceptors




acetic acid



trichloroacetic acid  
(herbicide, atmospheric  
breakdown product of  
chlorinated solvents)



benzoic acid (food  
preservative,  
additive, chemical  
intermediate)



(R,S)-2-(4-chloro-2-methyl  
phenyl) - propionic acid  
((R,S)-mecoprop; herbicide)


# Oxygen-containing functional groups

---


- **Ester groups**

- $R_1\text{-COO-}R_2$ ; -OH of a carboxylic acid is replaced by a -OR group
- Act only as a H-acceptor (smaller impact on a compound's water solubility)


ex) phthalates: often used as plasticizers



ethylacetate  
(acetic acid ethyl  
ester; solvent)



Preventol® B2  
(roof protection agent)













phthalates  
( $R_1, R_2 = C_1$  to  $C_{10}$ ;  
plasticizers)

# Nitrogen-containing functional groups

**Table 2.5** Some Important Nitrogen-Containing Functional Groups Present in Anthropogenic Organic Compounds

#4

Group	Name (oxidation state of nitrogen)	Group	Name (oxidation state of nitrogen)
	ammonium (-III)	$R_1-NH-NH-R_2$	hydrazo (-II)
	amino <sup>a</sup> (-III) (amine)		azo (-I)
	carboxylic acid amide <sup>a</sup> (-III)		hydroxyl-amine (-I)
$R-C\equiv N$	cyano, nitrilo (-III)		nitroso (+I)
	urea (-III)		nitro (+III)
	carbamate (-III)		nitrate (+V) (nitrate)



<sup>a</sup>Primary if  $R_2 = R_3 = H$ ; secondary if  $R_2 = H$  and  $R_3 \neq H$ ; tertiary if  $R_2 \neq H$  and  $R_3 \neq H$ .

# Nitrogen-containing functional groups


- **Amine groups**

- Types: primary/secondary/tertiary
- Natural/synthetic compounds
  - Natural example: amino acids
  - Synthetic example: anilines (intermediate for synthesis of dyes, pharmaceuticals, pesticides, antioxidants, ...), atrazine (pesticide)
- Acts as both H-acceptors and donors
  - H-acceptors: to a lesser extent
  - H-donors: only for primary and secondary amines
- Slightly basic: acquire a proton in an aqueous solution to form a cationic ammonium species

Amino Acid Structure



aniline and substituted  
anilines (chemical  
intermediates)



atrazine  
(a triazine herbicide)

#6




# Nitrogen-containing functional groups


---

- **Nitro groups**


- Widely used in the chemical industry
  - Explosives (ex: TNT), agrochemicals (ex: DNOC), dyes (ex: Dispersive Blue 79)
- Strong electron-withdrawing characteristics
  - Significantly affect the electron distribution in a molecule
  - Significantly affect the chemical properties of the compound
- Explosives: multiple nitro groups in the molecules
  - Nitro group as built-in oxidant
  - Very fast oxidation of the molecule



#2 2,4,6-trinitrotoluene  
(TNT, explosive)



2,4-dinitro-o-cresol  
(DNOC, herbicide)




Dispersive Blue 79  
(textile dye)


# Dissociation of nitrophenols

---

Compound	pK <sub>a</sub>
phenol	9.95
2-nitrophenol (ortho)	7.17
3-nitrophenol (meta)	8.28



**ortho-Nitrophenol or  
2-Nitrophenol**



**meta-Nitrophenol or  
3-Nitrophenol**

#7



# Sulfur-containing functional groups

**Table 2.6** Some Important Sulfur-Containing Functional groups Present in Anthropogenic Organic Compounds

#9

Group	Name (oxidation state of sulfur)	Group	Name (oxidation state of sulfur)
$R-SH$	thiol, mercaptan (-II)	$\begin{array}{c} O \\    \\ R-S-OH \\    \\ O \end{array}$	sulfonic acid (+IV)
$R_1-S-R_2$	thioether, sulfide (-II)	$\begin{array}{c} O \\    \\ R_1-S-O-R_2 \\    \\ O \end{array}$	sulfonic acid ester (+IV)
$\begin{array}{c} S \\    \\ R_1-C-R_2 \end{array}$	thiocarbonyl (-II)	$\begin{array}{c} O \\    \\ R_1-S-N \begin{array}{l} R_2 \\ R_3 \end{array} \\    \\ O \end{array}$	sulfonic acid amide, sulfonamide (+IV)
$R_1-S-S-R_2$	disulfide (-I)	$\begin{array}{c} O \\    \\ R_1-O-S-O-R_2 \\    \\ O \end{array}$	sulfuric acid ester, sulfate (+VI)
$\begin{array}{c} O \\    \\ R_1-S-R_2 \end{array}$	sulfoxide (0)		
$\begin{array}{c} O \\    \\ R_1-S-R_2 \\    \\ O \end{array}$	sulfone (+II)		

# References

---

- #1, #2, #4, #6, #9) Schwarzenbach, R., Gschwend, P. M., Imboden, D. M. (2003) *Environmental Organic Chemistry*, 2nd ed., John Wiley & Sons, p. 38, p. 40, p. 43, p. 44, p. 46.
- #3) [https://commons.wikimedia.org/wiki/File:Resonance\\_stabilization\\_of\\_carboxylic\\_acids.png](https://commons.wikimedia.org/wiki/File:Resonance_stabilization_of_carboxylic_acids.png)
- #5) <http://www.nutrientsreview.com/proteins/amino-acids>
- #7) <https://www.meritnation.com/ask-answer/question/explain-the-structure-of-nitrophenol/chemical-bonding-and-molecular-structure/3687611>
- #8) <https://sci-toys.com/attention/2006/04/octanitrocubane-most-powerful.html>

# W-Chem & O-Chem: Exercise

# Water solubility

---

**Q:** Among ethane ( $\text{C}_2\text{H}_6$ ), ethanol ( $\text{C}_2\text{H}_5\text{OH}$ ), hexachloroethane ( $\text{C}_2\text{Cl}_6$ ), which one will be the most soluble in water and which one the least? Describe your rationale.

# COD & TOC

---

**Q:** Compare the theoretical COD/TOC ratios for the following compounds.

acetic acid,  $\text{CH}_3\text{COOH}$

2-butanol,  $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$

glyceraldehyde,  $\text{CH}_2(\text{OH})\text{CH}(\text{OH})\text{CHO}$





2-chloropropane,  $\text{CH}_3\text{CHClCH}_3$



# Isomers

---

**Q:** A chemical structure (constitution) that has at least one pair of enantiomers is called as “chiral”. For each of chemical structures shown below, determine if it is chiral. If so, provide the total number of enantiomers.



# EDGs & EWGs

---

**Q:** Among the following molecules, which one will have the highest  $pK_a$  and which one the lowest? Describe your rationale.

- i) phenol
- ii) p-cresol (systematic name: 4-methylphenol)
- iii) 2-nitrophenol
- iv) 3-nitrophenol