Advanced Thermodynamics (M2794.007900)

Chapter 10

The Third Law of Thermodynamics

Min Soo Kim Seoul National University



- The third law of thermodynamics is concerned with the behavior of systems in equilibrium as their temperature approaches zero
- The definition of entropy given by

$$S = \int_0^T \frac{dQ_r}{T} + S_0 \tag{10.1}$$

is incomplete because of the undetermined additive constant S_0 , the entropy at absolute zero.

• In this chapter we shall introduce a principle that will enable us to determine S_0 .



• Gibbs-Helmholtz equation

$$G = H + T\left(\frac{\partial G}{\partial T}\right)_P \tag{10.2}$$

• If this relation is applied to the initial and final states of a system undergoing an isothermal process, it takes the form

$$\Delta G = \Delta H + T \left[\frac{\partial (\Delta G)}{\partial T}\right]_P \tag{10.3}$$

• This shows that the change in enthalpy and the change in the Gibbs function are equal at T = 0 for an isobaric process.

$$\lim_{T \to 0} \left[\frac{\partial (\Delta G)}{\partial T}\right]_P = 0 \qquad \lim_{T \to 0} \left[\frac{\partial (\Delta H)}{\partial T}\right]_P = 0 \tag{10.4}$$



• Equation (10.4) is illustrated in Figure 10.1.




Figure 10.1 Variation of ΔG and ΔH in the vicinity of absolute zero.



• We may write the equation (10.4) as

$$\lim_{T \to 0} \left[\frac{\partial (G_2 - G_1)}{\partial T}\right]_P = \lim_{T \to 0} \left[\left(\frac{\partial G_2}{\partial T}\right)_P - \left(\frac{\partial G_1}{\partial T}\right)_P\right] = 0$$
(10.5)

where the subscripts 1 and 2 refer to the initial and final states, respectively.

• From the reciprocity relation, we have the Nernst formulation of the third law.

$$\lim_{T \to 0} (S_1 - S_2) = 0 \tag{10.6}$$

All reactions in a liquid or solid in thermal equilibrium take place with no change of entropy in the neighborhood of absolute zero.

$$Cf.G = U - TS + PV \rightarrow dG = -SdT + VdP \rightarrow S = -\left(\frac{\partial G}{\partial T}\right)_{P}$$



• Planck extended Nernst's hypothesis by assuming that it holds for G_1 and G_2 separately.

$$\lim_{T \to 0} G(T) = \lim_{T \to 0} H(T)$$
(10.7)

$$\lim_{T \to 0} \left(\frac{\partial G}{\partial T}\right)_P = \lim_{T \to 0} \left(\frac{\partial H}{\partial T}\right)_P \tag{10.8}$$

• For convenience, we temporarily introduce a variable $\Phi \equiv G - H$. Equation (10.2) then becomes

$$T(\frac{\partial G}{\partial T})_P - \Phi = 0 \tag{10.9}$$

• Adding the term $-T(\partial H/\partial T)_P$ to both sides of Equation (10.9), we get

$$T(\frac{\partial \Phi}{\partial T})_P - \Phi = -T(\frac{\partial H}{\partial T})_P$$
(10.10)



• By L'Hopital's rule,

$$\lim_{T \to 0} \left(\frac{\partial H}{\partial T}\right)_P = 0 \tag{10.11}$$

• Finally, by Equation (10.8), we obtain the result

$$\lim_{T \to 0} \left(\frac{\partial G}{\partial T}\right)_P = 0 \tag{10.12}$$

• Since $\left(\frac{\partial G}{\partial T}\right)_P = -S$, it follows the Planck's statement of the third law.

$$\lim_{T \to 0} S = 0 \tag{10.13}$$

The entropy of a true equilibrium state of a system at absolute zero is zero

